

from 40.5-45.6 mm (Table 22), and there was a significant difference in depth between Regions 2 and 4 (Tables 23-24).

Quadrula pustulosa shell interval modes ranged from 30-39.9 to 40-49.9 mm in the four regions (Tables 18-21). The mean depth for this species ranged from 35.4-41.2 mm (Table 22), and there were significant differences in depth between Regions 1-2, 2-3 and 2-4 (Tables 23-24).

Discussion

Since this river is not commercially harvested, no selected size class of mussels is removed that would skew distribution of size classes in the community. Without the biased removal of legal size and larger individuals, the communities in the Ouachita River are more natural than those of other rivers where commercial harvesting occurs.

Species exhibiting differences in growth modes had the highest mode in the unchannelized Region 1. An alternate hypothesis suggests that mean size decreases as densities increase and resources decrease (Christian, 1995). Survey results suggest that greater intra- and interspecific competition for resources may affect observed size differences. Region 2 contained the highest estimated CNSC of all regions in the study area, and only *Fusconaia ebena* ($n=171$) showed a decrease in interval depth mode from Region 1 to Region 2 ($n=932$). Therefore, these data do not tend to support the hypothesis set forth by Christian (1995).

RECRUITMENT

Rust (1993) and Christian (1995) reported low recruitment within the rivers they studied. While few juvenile mussels were found in the St. Francis River, many juveniles were found in the Ouachita River. However, nearly all mussels collected in the Ouachita River were "dwarfed" when compared to those of other rivers and many "juveniles" may just be of a smaller age class and not true juveniles.

AGE DETERMINATIONS

St. Francis River

Results

Age analysis was conducted on 48 *Megalonaia nervosa* specimens ranging in depth from 79.8-130.8 mm. Age was determined to be from 8 to 41 years (Table 25, Figure 10). The correlation coefficient (r) was calculated to be 0.88, and r^2 was 0.77. Age at minimum legal harvestable size (92.2 mm), was predicted to be 9.9 years with standard error of prediction about the mean of 4.1 years.

Discussion

The r^2 value of 0.77 indicates a fairly strong correlation of the variability with only 23% of the variability unaccounted. Clinal variation and sexual dimorphism may explain some of the variability found (Oesch, 1984). The small sample size ($n=48$) may not provide enough data to precisely predict age at a certain depth. Experimental error may also provide some of the variability.

Table 25. Shell dimensions, dry mass and age for selected individuals of St. Francis River *Megalonaia nervosa*.


Number	Length	Depth	Width	Mass	Age
50	109.6	79.8	42.1	133	9
49	126.2	86.2	40.9	153	8
48	132.6	88.3	40.2	162	10
46	137.1	90.4	53.4	246	9
33	134.7	92.2	52.5	240	12
32	151.3	93.8	60.9	337	16
47	131.3	94.5	48.4	214	8
39	148.2	96.1	58.2	306	15
28	153.1	96.1	57.2	299	16
40	142.8	96.4	53.6	255	13
42	137.4	97.5	48.1	226	10
45	136.2	98.3	50.8	255	8
43	133.3	100.3	46.2	208	11
38	151.9	101.3	54.2	321	13
37	143.0	101.5	55.9	308	14
41	144.1	102.1	44.4	233	12
44	142.5	102.2	45.4	254	9
30	154.1	103.2	56.2	322	15
36	153.1	104.1	54.1	287	14
25	160.5	104.5	50.4	305	20
31	151.3	104.8	54.7	297	17
13	167.0	104.8	62.9	389	21
34	158.5	105.3	52.3	324	16
35	157.5	106.3	55.4	311	16
15	166.3	107.1	60.3	415	21
17	166.3	107.1	60.3	415	21
24	156.7	109.3	62.4	409	30
16	154.9	110.6	56.7	362	17
21	165.6	110.6	53.4	349	18
11	172.6	110.8	73.2	525	22
27	173.9	112.1	65.6	508	29
23	162.1	111.4	68.2	432	23
14	173.9	112.1	65.6	508	29
26	175.5	112.7	63.0	514	32
20	168.8	113.0	58.0	417	19
19	173.5	114.9	70.3	532	30
8	174.5	115.3	67.8	584	33
12	178.8	116.5	63.6	527	28
2	175.8	116.6	63.7	434	29
7	188.8	116.9	56.6	455	32
9	188.5	117.0	61.5	541	27
10	181.8	118.3	71.2	565	25
6	178.5	118.9	62.5	526	26
18	187.5	122.3	62.2	516	35
3	183.4	123.2	61.6	534	27
4	190.3	124.1	66.2	687	34
5	196.8	125.0	71.5	656	41

Table 25 con't.

Number	Length	Depth	Width	Mass	Age
1	205.5	130.8	71.0	647	37

Individuals 22 and 29 were broken during processing

Figure 10. Scatterplot of age versus depth for selected *Megalonaias nervosa* individuals, St. Francis River, Arkansas.



However, strict repetition was used to ensure duplicate results.

Correlating the age with commercial harvest size is important to further maintain a viable population, and thus a viable harvest. Woody and Holland-Bartels (1993) found *M. nervosa* of Pool 10 in the upper Mississippi river to be reproductively mature at age 8. Assuming that *Megalonaia* *nervosa* is sexually mature at this age in the St. Francis River, more protection may be needed to avoid overharvest of this species. However, more thorough studies need to be conducted to determine sexual maturity of this species in this river and to determine the minimum legal size from this age.

Ouachita River

Results

Fifty *Ambloema plicata* individuals ranging in depth from 19.3-64.1 mm were analyzed for age with counts ranging from 3-25 years (Table 26, Figure 11). The correlation coefficient (*r*) was calculated to be 0.72, and the *r*² was 0.52. Age at 69.9 mm (2.7") depth was predicted to be 24.4 years, with a standard error of prediction of 2.9 years.

Discussion

The *r*² value of 0.52 indicates a moderate correlation between depth and age and leaves 48% of the variability unexplained. No specimens aged were of legal harvestable size and all were collected from Region 1. Additionally, 70% of all specimens were less than 11 years old. Mussels tend to


Table 26. Shell dimensions, dry mass and age for individuals of selected Ouachita River *Amblema plicata*.

Number	Length	Depth	Width	Weight	Age
50	24.3	19.3	14.3	5	3
48	32.7	26.7	19.7	12	4
49	32.7	26.9	19.2	11	4
46	38.5	29.3	20.6	14	4
47	36.1	30.1	20.3	13	3
45	40.9	33.9	22.1	19	5
43	41.5	34.2	25.9	19	5
44	44.4	35.3	24.7	21	5
42	43.8	37.5	24.9	23	7
38	45.8	38.4	27.6	27	7
41	43.3	38.5	27.3	27	5
40	46.4	39.0	26.0	25	6
35	51.4	39.2	27.7	33	6
39	45.1	40.0	25.9	24	7
37	52.6	40.4	26.1	27	7
32	50.6	40.7	28.9	31	6
36	50.9	41.0	27.3	29	7
30	54.4	42.0	27.7	31	8
34	52.3	42.0	27.6	30	7
21	55.4	42.3	30.5	34	8
22	54.3	42.7	29.9	37	8
31	53.3	43.3	32.4	40	7
33	53.3	43.3	32.4	40	7
19	55.7	45.2	30.2	36	8
23	56.9	45.7	31.3	42	8
29	58.0	46.4	30.8	47	9
25	61.4	46.4	29.3	37	9
20	57.8	46.8	33.1	44	7
27	61.3	47.4	32.4	49	10
24	56.4	48.3	30.8	39	7
28	59.4	48.3	33.1	52	8
18	57.2	48.3	28.8	37	10
26	63.4	48.4	30.5	40	6
17	64.1	49.9	31.5	46	8
16	62.3	50.5	29.5	44	7
13	77.1	52.5	36.0	65	16
14	73.6	53.3	32.6	68	18
15	69.9	55.2	34.4	61	15
12	66.1	55.3	31.9	54	13
11	76.2	55.9	39.2	93	13
6	74.4	56.0	36.9	81	18
9	79.1	56.7	36.4	79	15
10	75.4	58.9	33.5	64	18
4	75.4	59.0	41.3	103	22
5	81.7	60.4	41.2	99	25
7	83.3	60.4	39.4	105	19
3	78.5	60.9	45.4	109	16

Table 26 con't.

Number	Length	Depth	Width	Weight	Age
2	87.4	63.2	42.0	123	21
1	88.5	63.6	35.1	75	25
8	82.2	64.1	39.1	106	13

Figure 11. Scatterplot of age versus depth for selected *Ambloema plicata* individuals, Ouachita River, Arkansas.



grow less as they age which may explain some of the variability (Figures 10 and 11).

PERCENT LEGAL HARVESTABLE MUSSELS

Knowing the percentage of legal harvestable mussels is necessary to determine the current status of mussel communities. It will also help to maintain a viable commercial industry for the future.

St. Francis River

Results

Within the St. Francis River, 41.3% of all mussels collected were of legal harvestable dimension. Mbeds contained 40.7% legal dimension individuals and mbeds contained 42.9% legal individuals (Table 27). Percent legal harvestable mussels (PLHM) for individual species is summarized in Table 28. PLHM data for M- and mbeds are summarized in Appendixes 1.1 and 1.2, respectively.

The highest mean PLHM was found in Region 1 and decreased through Regions 2 and 3 (Table 28). Bed 1m had the highest PLHM (69.6), while Bed 9m had the lowest (15.2).

Discussion

The significant current influence on PLHM in the St. Francis River is harvest pressure. Between 1990 and 1995, 792,516 kg of mussel shell were harvested from this river (1992 not reported, AGFC, 1993). More tonnage of shell material was removed from the St. Francis River from 1990, 1993 and 1994 than any other commercially harvested area

Table 27. Percent legal harvestable mussels, commercial species, in the St. Francis River, Arkansas.

	Mean	Segment 1	2	3
Mbeds	40.7 (467)	*	41.5 (406)	27.9 (61)
mbeds	42.9 (296)	51.9 (165)	46.7 (26)	15.2 (105)
Total	41.3 (763)	51.9 (165)	41.6 (432)	19.9 (166)

() = Sum of individuals

* = No major beds defined in this segment

Table 28. Percent legal harvestable mussels, by species, in the St. Francis River, Arkansas, 1993.

Species	Bed		
	Major	Minor	Mean
<i>Amblema plicata</i>	77.2 (338)	88.0 (132)	79.9 (470)
<i>Ellipsaria lineolata</i>	66.7 (2)	0.0 (0)	66.7 (2)
<i>Fusconaia ebena</i>	35.3 (6)	100.0 (1)	38.9 (7)
<i>Fusconaia flava</i>	31.3 (5)	50.0 (4)	37.5 (9)
<i>Megalonaia nervosa</i>	86.8 (33)	80.0 (8)	85.4 (41)
<i>Obliquaria reflexa</i>	7.7 (1)	20.0 (1)	11.1 (2)
<i>Plectomerus dombeyanus</i>	66.7 (2)	100.0 (1)	75.0 (3)
<i>Pleurobema coccineum</i>	40.0 (6)	0.0 (0)	40.0 (6)
<i>Pleurobema pyramidatum</i>	5.7 (2)	85.7 (6)	19.0 (8)
<i>Potamilus purpuratus</i>	100.0 (10)	85.7 (6)	94.1 (16)
<i>Quadrula metanevra</i>	100.0 (5)	0.0 (0)	100.0 (5)
<i>Quadrula nodulata</i>	14.3 (2)	16.7 (1)	15.0 (3)
<i>Quadrula pustulosa</i>	1.6 (4)	16.3 (21)	65.8 (25)
<i>Quadrula quadrula</i>	2.9 (4)	4.5 (5)	3.6 (9)
<i>Tritogonia verrucosa</i>	25.0 (2)	100.0 (11)	68.4 (13)
Mean	40.6 (422)	40.8 (197)	40.7 (619)

() = Sum Individuals

reported in the state. In 1995, only Macon Bayou and Fourche LaFave River produced more commercial shell material than the St. Francis River (AGFC, 1996).

Conversations with local shell fishermen attribute the shallow depth, allowing easy access to mussels in the St. Francis River, as the major reason for the heavy harvest. Often, mussels are collected by wading and feeling for mussels in the substrate. This is a low cost alternative to the more expensive SCUBA required to harvest in deeper rivers.

Ouachita River

Results

The Ouachita River produced only 0.3 PLHM, all were found in Mbeds. Region 2 contained the greatest number of legal individuals (43) while Region 4 contained the fewest (3). Of notable interest, these two regions contained the lowest mean density per region, 25.5 and 24.1 mussels/m², respectively. Values for PLHM for M- and mbeds are summarized in Table 29. Individual PLHM values for species are found in Table 30 and individual M- and mbed PLHM's are found in Appendices 1.3 and 1.4, respectively.

The greatest PLHM was 1.0 (25M) found in Region 2. Forty-two (68.9%) beds contained no legal size mussels.

Discussion

Commercial harvesting is permitted from the Clark-Ouachita County Line to Felsenthal National Refuge and from Felsenthal Lock and Dam to Louisiana. No harvesting occurs

Table 29. Percent legal harvestable mussels, commercial species, in the Ouachita River, Arkansas.

	Region			
	Mean	1	2	3
Major Beds	0.3 (21,371)	0.1 (4,036)	0.5 (8,147)	0.6 (956)
Minor Beds	0.0 (2,006)	0.0 (885)	0.0 (63)	0.0 (421)
Total	0.3 (23,463)	0.08 (4,921)	0.05 (8,210)	0.4 (1,377)
				0.03 (8,955)

() = Sum of individuals

Table 30. Percent legal harvestable mussels, by species, in the Ouachita River, Arkansas, 1993-1995.

Species	Bed		
	Major	Minor	Mean
<i>Actinonaias ligamentina</i>	3.0 (4)	0.0(0)	2.4 (4)
<i>Amblema plicata</i>	0.3 (10)	0.0(0)	0.2 (10)
<i>Fusconaia ebena</i>	0.4 (14)	0.0(0)	0.4 (14)
<i>Fusconaia flava</i>	0.04 (1)	0.0(0)	0.04 (1)
<i>Megalonaia nervosa</i>	1.1 (8)	0.0(0)	1.1 (8)
<i>Pleurobema pyramidatum</i>	0.09 (3)	0.0(0)	0.08 (3)
<i>Plectomerus dombeyanus</i>	0.6 (6)	0.0(0)	0.6 (6)
<i>Potamilus purpuratus</i>	7.4 (9)	0.0(0)	7.0 (9)
<i>Quadrula metanevra</i>	0.3 (1)	0.0(0)	0.2 (1)
<i>Tritogonia verrucosa</i>	0.4 (1)	0.0(0)	0.3 (1)
Mean	0.3 (57)	0.0(0)	0.3 (57)

() = Sum of individuals

because there are few legal sized mussels encountered.

A large proportion of the collected valves had umbonal erosion exceeding 1.0 cm. Umboonal erosion has been related to substrate particle size by Kesler and Bailey (1993). Their study found that mussels inhabiting sand substrate had more erosion than those inhabiting silt substrate. However, regardless of substrate, older individuals were more eroded than younger individuals. Sand was often found in the interstices of the gravel substrate and may be the cause of the shell erosion found in this river.

CHAPTER V

MANAGEMENT RECOMMENDATIONS

The goal of this research was to map and record community parameters for commercial size mussel beds in order to provide baseline data to facilitate proper management of this resource in the future. With mapping of these beds, habitat loss due to dredging and river side construction in these areas should be alleviated. Locations of endangered species will also assist in preserving these habitats.

Historically, 63 known mussel beds were located in the St. Francis River. However, only 10 of these were mapped and sampled during this survey. The other 53 sites either did not meet bed criteria or were absent. More protection for this river is necessary if the communities present are expected to survive. The Refuge established from the I-40 bridge downstream to the Ark. Hwy. 50 bridge at Madison in St. Francis County restricting commercial harvest will supply some of this protection. However, none of the mapped beds fall within the boundaries of this refuge. An additional refuge should be placed below Parkin (RM 75.3-82.5), where 75% of all Mbeds were mapped. Sampling this area yielded 978 mussels (64.3% of total) representing 20 mussel species, more than any other region.

Another suggestion would be to close all harvest on the St. Francis River for several years and trace recruitment into areas affected by the restrictions. At the present time, the low number of juveniles of commercial species in surveyed areas is disconcerting to those managing this resource.

Heavy siltation in the St. Francis River needs to be addressed. Flocculent silt is unsuitable for mussels due to the decreased water movement in the vicinity of the animal (Oesch, 1984). Historically, this river was described as a clear stream with sand substrate (Meek, 1896). The presence of silt substrates encountered suggests erosion from the agriculture dominated watershed (ADPCE, 1987). Perhaps more "no till" planting could help reduce the amount of silt entering the river. Buffer zones between the fields and river would help filter sediments before entering the river as well as reduce bank erosion, thus saving the farmers' land.

Commercial harvesting should no longer be permitted in the Ouachita River. A viable mussel industry does not exist in this river, and permitting commercial harvest seems without merit. If a viable industry should come to be in the future, no shelling should be permitted upstream of Camden, Ark. The majority of endangered species encountered during this survey were found in the region extending from Clark County to Camden. This area is not currently maintained by the USACOE, nor is there any reason for maintenance.

Because of the potential threat of the zebra mussel,

every effort should be made to retard the dispersal of this exotic species. Posters and wash stations will serve to educate the public, but the movement into new habitats is inevitable. The ongoing plans are to slow the spread of this exotic species and buy time in hope some form of control will be found before irreversible damage to local fauna occurs.

No single management tool will sustain current mussel communities. Some streams in Arkansas will no longer support a viable community of organisms with the present degree of impact from pollution and channelization. It is my hope that the information established for these two surveyed rivers will stimulate restoration efforts before they become unsuitable for aquatic life.

LITERATURE CITED

- Ahlstedt, S.A. and J.J. Jenkinson. 1987. Distribution and abundance of *Potamilus capax* and other freshwater mussels in the St. Francis River System, Arkansas and Missouri. Final Report. USACOE. Memphis Dist. 67 pp. + append.
- Ahlstedt, S.A. and J.J. Jenkinson. 1991. Distribution and abundance of *Potamilus capax* and other freshwater mussels in the St. Francis River System, Arkansas and Missouri. Walkerana. 5(14):225-261.
- Allan, J.D. and A.S. Flecker. 1993. Biodiversity conservation in running waters. Bioscience. 43(1):32-42.
- ADPCE. 1979. Section 303(c) Basin Plan (Ouachita River Basin) 563 pp.
- ADPCE. 1987. Physical, chemical and biological characteristics of least-disturbed reference streams in Arkansas Ecoregions. Ecoregion Reference Streams. Vol. 1. 685 pp.
- AGFC. 1991. 1990 mussel harvest report. Fisheries Division, Little Rock. April 19. 6 pp.
- AGFC. 1992. 1991 mussel harvest report. Fisheries Division, Little Rock. March 18. 4 pp.
- AGFC. 1993. 1992 mussel harvest report. Fisheries Division, Little Rock. March 5. 11 pp.
- AGFC. 1994. 1993 mussel harvest report. Fisheries Division, Little Rock. May 12. 15 pp.
- AGFC. 1995. 1994 mussel harvest report. Fisheries Division, Little Rock. May 8. 15 pp.
- AGFC. 1996. 1995 mussel harvest report. Fisheries Division, Little Rock. May 7. 16 pp.
- Bates, J.M and S.D. Dennis. 1983. Mussel (naiad) survey-St. Francis, White, and Cache rivers, Arkansas and Missouri. U.S. Army Corps of Engineers, Memphis. Project # DACW66-78-C-0147. 88 pp. + appendices A-E.

- Call, R.E. 1895. A study of the Unionidae of Arkansas, with incidental reference to their distribution in the Mississippi Valley. Trans. Mo. Acad. Sci. St. Louis 7:1-65.
- Christian, A.D. 1995. Analysis of the commercial mussel beds in the Cache and White Rivers in Arkansas. Unpubl. M.S. thesis. Arkansas State University, State University, AR. 197 pp.
- Cox, G.W. 1980. A laboratory manual of general ecology. 5th Ed. William C. Brown Co. Publishers. Dubuque, Iowa. 237 pp.
- Cummings, K.S. and C.A. Mayer. 1992. Field guide to freshwater mussels of the midwest. Illinois Natural History Survey. Manual 5. 194 pp.
- Ecosearch, Inc. 1985. Mussel (naiad) study: St. Francis and White rivers Arkansas. U.S. Army Corps of Engineers, Memphis. Project # DAC-84-M-1666R. 29 pp.
- Gordon, M.E. 1980. Recent Mollusca of Arkansas with annotations to systematics and zoogeography. Proc. Ark. Acad. Sci. 34:58-62.
- Gordon, M.E. and J.B. Layzer. 1989. Mussels (BIVALVA:UNIONIDAE) of the Cumberland River: review of life histories and ecological relationships. U.S. Fish Wld. Serv. Bio. Rep. 89(15). 99 pp.
- Gordon, M.E., L.R. Kraemer and A.V. Brown. 1979. Unionaceae of Arkansas: Historical review, checklist, and observations on distribution patterns. Am. Mal. Bull. 1979:31-37.
- Haage, W.R., D.J. Berg, S.W. Garton and J.L. Farris. 1993. Reduced survival and fitness in native bivalves in response to fouling by the introduced zebra mussel (*Dreissena polymorpha*) in western Lake Erie. Can.J. Fish. Aquat. Sci. 50:13-19.
- Harris, J.L. 1986. Relocation of the fat pocketbook pearly mussel, *Proptera capax* (Green), in the St. Francis River at Madison, St. Francis County, Arkansas. Ark. Highway and Transport. Dept. and Ark. Game and Fish Comm., Little Rock. 14 pp.
- Harris, J.L. 1997. A population assessment of recolonization by the fat pocketbook mussel of dredged habitat in the St. Francis Floodway, Arkansas. Final Report. 14 pp + append.

- Harris, J.L. and M.E. Gordon. Undated. Arkansas mussels. Arkansas Game and Fish Commission. Little Rock, Ark. 32 pp.
- Harris, J.L. and M.E. Gordon. 1985. Survey of the Ouachita River in Arkadelphia, Arkansas for the endangered freshwater mussel *Lampsilis orbiculata* (Hildreth). Arkansas Louisiana Gas Company, Shreveport, LA. 8 pp.
- Harris, J.L. and M.E. Gordon. 1987. Distribution and status of rare and endangered mussels (Mollusca: Margaritiferidae, Unionidae) in Arkansas. Proc. Ark. Acad. Sci. 41:49-55.
- Harris, J.L., P. Rust, S.W. Chordas III and G.L. Harp. 1993. Distribution and population structure of freshwater mussels (Unionidae) in Lake Chicot, Arkansas. Proc. Ark. Acad. Sci. 47:38-43.
- Hesse, L.S. and W. Sheets. 1993. The Missouri River hydrosystem. Fisheries. 18(5):5-14.
- Holland-Bartels, L.E. 1990. Physical factors and their influence on the mussel fauna of a main channel border habitat of the upper Mississippi River. J.N. Am. Benthol. Soc. 9(4):327-335.
- Huebner, J.D., D.F. Malley and K. Donkersloot. 1990. Population ecology of the freshwater mussel *Anodonta grandis grandis* in a Precambrian Shield lake. Can. J. Zool. 68:1931-1941.
- Jenkinson, J.J. 1989. Relocation of *Potamilus capax* from a 4-mile reach of the St. Francis River Floodway in Arkansas. Final Report. USACOE. Memphis Dist. 17 pp. + append.
- Jenkinson, J.J and S.A. Ahlstedt. 1988. A search for additional populations of *Potamilus capax* in the St. Francis and Cache river watersheds, Arkansas and Missouri. Final Report. USACOE. Memphis Dist. 105 pp + append.
- Kesler, D.H. and R.C. Bailey. 1993. Density and ecomorphology of a freshwater mussel (*Elliptio complanata*, Bivalva: Unionidae) in a Rhode Island lake. J. N. Am. Benthol. Soc. 12(3):259-264.
- LORWG. 1993. A survey of the fish community in the Lower Ouachita River, Arkansas. Final Report. 47 pp.

- Ludyanskiy, M.L., D. McDonald and D. MacNeil. 1993. Impact of the zebra mussel, a bivalve invader. BioScience 43(8):533-541.
- Madison, J. 1985. Mississippi shell game. Audobon. 87(2): 46-48.
- Meek, S.E. 1896. A list of fishes and mollusks collected in Arkansas and Indian Territory in 1894. Bull. U.S. Fish. Comm. 15:341-349.
- Neves, R.J. and S.N. Moyer. 1988. Evaluation of techniques for age determination of freshwater mussels(Unionidae). Am. Mal. Bull. 6(2): 179-188.
- Oesch, R.D. 1984. Missouri naiades, a guide to the mussels of Missouri. Missouri Department of Conservation, Jefferson City. 270 pp.
- Parmalee, P.W. 1967. The freshwater mussels of Illinois. Illinois State Museum. Popular Sci. Series. Vol. 8 108 pp. in Oesch, R.D. 1984. Missouri naiades, a guide to the mussels of Missouri. Missouri Department of Conservation, Jefferson City. 270 pp.
- Posey, W.R. II, J.L. Harris and G.L. Harp. 1996. New distributional records for freshwater mussels in the Ouachita River, Arkansas. Proc. Ark. Acad. Sci. 50:96-98.
- Rust, P.J. 1993. Analysis of the commercial mussel beds in the Black, Spring, Strawberry and Current rivers in Arkansas. Unpubl. M.S. thesis, Arkansas State University, State University, AR. 125 pp.
- Sampford, M.R. 1962. An introduction to sampling theory with applications to agriculture. Oliver & Boyd Ltd., Edinburgh.
- Sheldon, A.L. 1968. Species diversity and longitudinal succession in stream fishes. Ecology. 49(2):193-198.
- Sokal, R.R. and F.J. Rohlf. 1981. Biometry. Second ed. W.H. Freeman and Company. San Francisco. 859 pp.
- Stansbery, D.H. and C.B. Stein. 1982. The unionid mollusks of the Lower St. Francis River in Arkansas. U.S. Fish and Wildlife Serv. Final Report. Contract - P.O. #98210-0884. 24 pp + append.
- USACOE. 1987. Ouachita and Black Rivers. Map and information pamphlet. 12 pp.

- USGS 1972. Drainage area of streams in Arkansas. United States Geological Survey. 32pp.
- USGS 1979. Drainage areas of streams in Arkansas. United States Geological Survey. 87 pp.
- Watters, G.T. 1993. Mussel diversity as a function of drainage area and fish diversity: management implications. Pages 113-116 in K.S. Cummings, A.C. Buchanan, and L.M. Koch, eds. Conservation and management of freshwater mussels. Proceedings of a UMRCC symposium, 12-14 October 1992, St. Louis, Missouri. Upper Mississippi River Conservation Committee, Rock Island, Illinois. 189 pp.
- Wheeler, 1918. The Mollusca of Clark County, Arkansas. *Nautilus*. 31(4):109-125.
- Williams, J.D., M.L. Warren, Jr., K.S. Cummings, J.L. Harris, and R.J. Neves. 1993. Conservation status of freshwater mussels of the United States and Canada. *Am. Fish. Soc.* 18(9):6-22.
- Witte, R.S. 1993. Statistics. 4th Ed. Harcourt Brace Jonanovich Publishers. Orlando, FL. 551 pp.
- Woody, C.A. and L.E. Holland-Bartels. 1993. Reproductive characteristics of a population of the washboard mussel, *Megalonaia nervosa* (Rafinesque 1820) in the Upper Mississippi River. *Freshwater Ecology*. 8(1):54-66.

APPENDIX 1.1 Physical parameters, species composition,
 percent legal harvestable mussels, population
 estimates and community estimates for major
 beds of the St. Francis River, Arkansas,
 1994.

BED: 6M
 Location: Mile 82.5
 AR: Cross County
 Stratum Size: 200m x 3m = 600m²
 Total Samples: 10
 Substrate: Sand, soft clay
 Minimum-Maximum density (#/m²): 2.0-45.0
 Mean density #/m² (Standard deviation): 23.2 (15.0)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	61	26.3	77.7	3660±2245
<i>Arcidens confragosus</i>	3	1.3	*NV	180±287
<i>Ellipsaria lineolata</i>	2	0.9	100.0	120±179
<i>Fusconaia ebena</i>	13	5.6	30.8	780±1135
<i>Fusconaia flava</i>	9	3.9	55.6	540±372
<i>Megalonaia nervosa</i>	11	4.7	91.0	660±678
<i>Obliquaria reflexa</i>	4	1.7	25.0	240±359
<i>Pleurobema pyramidatum</i>	9	3.9	0.0	540±813
<i>Potamilus purpuratus</i>	4	1.7	100.0	240±219
<i>Quadrula metanevra</i>	3	1.3	100.0	180±287
<i>Quadrula nodulata</i>	6	2.6	33.3	360±574
<i>Quadrula pustulosa</i>	75	32.3	4.0	4500±2357
<i>Quadrula quadrula</i>	26	11.2	11.5	560±901
<i>Tritogonia verrucosa</i>	3	1.3	0.0	180±287
<i>Truncilla donaciformis</i>	1	0.4	NV	60±134
<i>Truncilla truncata</i>	2	0.9	NV	120±269
Totals	232	100.0	36.6	14100±6387

NV = No Commercial Value

Appendix 1.1 cont'd

BED: 7M

Location: Mile 75.8

AR: Cross County

Stratum Size: 150m x 4m = 600m²

Total Samples: 10

Substrate: Sand

Minimum-Maximum density (#/m²): 1.0-79.0Mean density #/m² (Standard deviation): 24.4 (21.7)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	123	50.4	83.7	1200±2753
<i>Arcidens confragosus</i>	6	2.5	*NV	360±457
<i>Fusconaia ebena</i>	1	0.4	100.0	60±134
<i>Fusconaia flava</i>	4	1.6	0.0	240±411
<i>Leptodea fragilis</i>	9	3.7	NV	540±616
<i>Megalonaia nervosa</i>	5	2.0	80.0	300±540
<i>Obliquaria reflexa</i>	6	2.5	0.0	360±457
<i>Pleurobema coccineum</i>	1	0.4	100.0	60±134
<i>Potamilus purpuratus</i>	2	0.8	100.0	120±179
<i>Quadrula metanevra</i>	1	0.4	100.0	60±134
<i>Quadrula nodulata</i>	2	0.8	0.0	120±269
<i>Quadrula pustulosa</i>	56	23.0	1.8	3360±3233
<i>Quadrula quadrula</i>	24	9.8	0.0	1440±901
<i>Tritogonia verrucosa</i>	2	0.8	50.0	120±269
<i>Truncilla donaciformis</i>	1	0.4	NV	60±134
<i>Truncilla truncata</i>	1	0.4	NV	60±134
Totals	244	99.9	46.7	21960±13915

* NV = No Commercial Value

Appendix 1.1 cont'd

BED: 8M
 Location: Mile 75.3
 AR: Cross County
 Stratum Size (n samples): 1.) 100m x 3m = 300m² (5)
 2.) 800m x 3m = 2400m² (19)
 Total Samples: 24
 Substrate: 1.) Soft clay
 2.) Sand, soft clay
 Minimum-Maximum density (#/m²): 1.) 7.0-65.0
 2.) 3.0-47.0
 Mean density #/m² (Standard deviation): 1.) 30.6 (24.8)
 2.) 18.3 (14.1)
 Total: 20.9 (17.0)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	246	49.0	73.2	30079±11865
<i>Ellipsaria lineolata</i>	1	0.2	0.0	60±123
<i>Fusconaia ebena</i>	2	0.4	0.0	120±245
<i>Fusconaia flava</i>	3	0.6	0.0	180±368
<i>Lasmigona complanata</i>	2	0.4	*NV	120±150
<i>Leptodea fragilis</i>	9	1.8	NV	1004±691
<i>Megalonaia nervosa</i>	15	3.0	80.0	1828±1595
<i>Obliquaria reflexa</i>	3	0.6	0.0	246±349
<i>Plectomerus dombeyanus</i>	2	0.4	100.0	186±283
<i>Pleurobema coccineum</i>	14	2.8	35.7	840±1051
<i>Pleurobema pyramidatum</i>	26	5.2	7.7	1626±1311
<i>Potamilus purpuratus</i>	4	0.8	100.0	439±462
<i>Quadrula metanevra</i>	1	0.2	100.0	126±258
<i>Quadrula nodulata</i>	5	1.0	0.0	499±449
<i>Quadrula pustulosa</i>	120	23.9	0.0	1511±5931
<i>Quadrula quadrula</i>	44	8.8	2.3	3966±1894
<i>Tritogonia verrucosa</i>	3	0.6	33.3	246±345
<i>Truncilla truncata</i>	2	0.4	NV	120±150
Totals	502	100.1	41.4	53198±17145

* NV = No Commercial Value

Appendix 1.1 cont'd

BED: 10M

Location: Mile 6.0

AR: Lee County

Stratum Size: 150m x 6m = 900m²

Total Samples: 10

Substrate: Soft clay

Minimum-Maximum density (#/m²): 0.0-13.0Mean density #/m² (Standard deviation): 6.1 (4.1)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	8	13.1	100.0	720±526
<i>Fusconaia ebena</i>	1	1.6	100.0	90±202
<i>Lasmigona complanata</i>	1	1.6	*NV	90±202
<i>Megalonaia nervosa</i>	7	11.5	100.0	630±526
<i>Plectomerus dombeyanus</i>	1	1.6	100.0	90±202
<i>Quadrula nodulata</i>	1	1.6	0.0	90±202
<i>Quadrula quadrula</i>	42	68.9	0.0	3780±2252
Totals	61	99.9	27.9	5400±2646

* NV = No Commercial Value

APPENDIX 1.2 Physical parameters, species composition, percent legal harvestable mussels and mean densities for minor beds of the St. Francis River, Arkansas, 1994.

BED: 1m

Location: Mile 123.3

AR: Poinsett County

Stratum Size: 200m x 2m = 400m²

Total Samples: 5

Substrate: Sand, soft clay

Minimum-Maximum density (#/m²): 4.0-14.0

Mean density #/m² (Standard deviation): 9.2 (3.6)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	30	65.2	93.3	6.0
<i>Arcidens confragosus</i>	1	2.2	*NV	0.2
<i>Lasmigona complanata</i>	2	4.3	NV	0.4
<i>Leptodea fragilis</i>	1	2.2	NV	0.2
<i>Potamilus purpuratus</i>	2	4.3	100.0	0.4
<i>Pyganodon grandis</i>	1	2.2	NV	0.2
<i>Quadrula pustulosa</i>	6	13.0	33.3	1.2
<i>Quadrula quadrula</i>	1	2.2	0.0	0.2
<i>Tritogonia verrucosa</i>	1	2.2	0.0	0.2
<i>Truncilla truncata</i>	1	2.2	NV	0.2
Totals	46	100.0	69.6	9.0

* NV = No Commercial Value

Appendix 1.2 cont'd.

BED: 2m
 Location: Mile 123.0
 AR: Poinsett County
 Stratum Size: 80m x 4m = 320m²
 Total Samples: 5
 Substrate: Sand, soft clay
 Minimum-Maximum density (#/m²): 4.0-21.0
 Mean density #/m² (Standard deviation): 10.8 (6.2)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	10	18.5	90.0	2.0
<i>Fusconaia ebena</i>	1	1.8	100.0	0.2
<i>Fusconaia flava</i>	2	3.7	100.0	0.4
<i>Lampsilis teres</i>	2	3.7	100.0	0.4
<i>Leptodea fragilis</i>	15	27.8	NV*	3.0
<i>Megalonaia nervosa</i>	1	1.8	100.0	0.2
<i>Potamilus purpuratus</i>	2	3.7	50.0	0.4
<i>Pyganodon grandis</i>	1	1.8	NV	0.2
<i>Quadrula pustulosa</i>	9	16.7	22.2	1.2
<i>Quadrula quadrula</i>	7	13.0	14.3	1.4
<i>Tritogonia verrucosa</i>	4	7.4	100.0	0.7
Totals	54	99.9	42.6	10.8

* NV = No Commercial Value

Appendix 1.2 cont'd.

BED: 3m

Location: Mile 121.6

AR: Poinsett County

Stratum Size: 100m x 2m = 200m²

Total Samples: 5

Substrate: Sand, clay, broken glass

Minimum-Maximum density (#/m²): 1.0-68.0Mean density #/m² (Standard deviation): 21.8 (26.9)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	26	23.8	84.6	5.2
<i>Fusconaia flava</i>	5	4.6	20.0	1.0
<i>Lasmigona complanata</i>	1	0.9	NV*	0.2
<i>Leptodea fragilis</i>	3	2.8	NV	0.6
<i>Megalonaia nervosa</i>	3	2.8	100.0	0.6
<i>Obliquaria reflexa</i>	1	0.9	100.0	0.2
<i>Pleurobema pyramidatum</i>	3	2.8	100.0	0.6
<i>Potamilus purpuratus</i>	1	0.9	100.0	0.2
<i>Quadrula nodulata</i>	1	0.9	0.0	0.2
<i>Quadrula pustulosa</i>	61	56.0	14.8	12.2
<i>Quadrula quadrula</i>	2	1.8	50.0	0.4
<i>Tritogonia verrucosa</i>	2	1.8	100.0	0.4
Totals	109	100.0	39.4	21.8

* NV = No Commercial Value

Appendix 1.2 cont'd.

BED: 4m

Location: Mile 120.6

AR: Poinsett County

Stratum Size: 200m x 2m = 200m²

Total Samples: 5

Substrate: Sand, gravel and clay

Minimum-Maximum density (#/m²): 6.0-46.0Mean density #/m² (Standard deviation): 21.8 (15.0)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	52	47.7	98.1	10.4
<i>Leptodea fragilis</i>	7	6.4	NV*	1.4
<i>Megalonaia nervosa</i>	1	0.9	100.0	0.2
<i>Obliquaria reflexa</i>	2	1.8	0.0	0.2
<i>Pleurobema pyramidatum</i>	2	1.8	100.0	0.4
<i>Potamilus purpuratus</i>	2	1.8	100.0	0.4
<i>Quadrula nodulata</i>	2	1.8	50.0	0.4
<i>Quadrula pustulosa</i>	29	26.6	27.6	5.8
<i>Quadrula quadrula</i>	9	8.3	0.0	1.8
<i>Tritogonia verrucosa</i>	3	2.8	66.7	0.6
Totals	109	99.9	61.5	21.8

* NV = No Commercial Value

Appendix 1.2 cont'd.

BED: 5m

Location: Mile 85.0

AR: Cross County

Stratum Size: 200m x 2m = 400m²

Total Samples: 5

Substrate: Sand, clay

Minimum-Maximum density (#/m²): 7.0-18.0Mean density #/m² (Standard deviation): 12.0 (5.3)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	24	40.0	91.7	4.8
<i>Fusconaia flava</i>	1	1.7	100.0	0.2
<i>Leptodea fragilis</i>	1	1.7	NV*	0.2
<i>Megalonaia nervosa</i>	2	3.3	100.0	0.4
<i>Plectomerus dombeyanus</i>	1	1.7	100.0	0.2
<i>Pleurobema pyramidatum</i>	2	3.3	50.0	0.4
<i>Quadrula nodulata</i>	2	3.3	0.0	0.4
<i>Quadrula pustulosa</i>	22	36.7	0.0	4.4
<i>Quadrula quadrula</i>	4	6.7	0.0	0.8
<i>Tritogonia verrucosa</i>	1	1.7	100.0	0.2
Totals	60	100.1	46.7	12.0

* NV = No Commercial Value

Appendix 1.2 cont'd.

BED: 9m

Location: Mile 11.45

AR: Lee County

Stratum Size: 60m x 4m = 240m²

Total Samples: 5

Substrate: Stiff clay, gravel

Minimum-Maximum density (#/m²): 1.0-62.0Mean density #/m² (Standard deviation): 21.4 (24.6)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	8	7.6	100.0	1.6
<i>Megalonaia nervosa</i>	3	2.9	100.0	0.6
<i>Obliquaria reflexa</i>	2	0.8	0.0	0.8
<i>Quadrula nodulata</i>	1	0.9	0.0	0.2
<i>Quadrula pustulosa</i>	2	1.9	0.0	0.4
<i>Quadrula quadrula</i>	89	84.8	3.4	17.8
<i>Tritogonia verrucosa</i>	2	1.8	100.0	0.4
Totals	107	100.0	13.3	21.0

* NV = No Commercial Value

APPENDIX 1.3 Physical parameters, species composition, percent legal harvestable mussels, population estimates and community estimates for major beds of the Ouachita River, Arkansas, 1993-95.

BED: 1M

Location: Mile 376.2

AR: Ouachita County

Stratum Size: 200m x 6m = 1200m²

Total Samples: 12

Substrate: Gravel

Minimum-Maximum density (#/m²): 5.0-42.0

Mean density #/m² (Standard deviation): 19.6 (12.1)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	9	3.8	0.0	900±657
<i>Amblema plicata</i>	32	13.6	0.0	3200±2177
<i>Cyprogenia aberti</i>	1	0.4	0.0	100±218
<i>Ellipsaria lineolata</i>	2	0.9	0.0	200±295
<i>Elliptio dilatata</i>	32	13.6	0.0	3200±2177
<i>Fusconaia ebena</i>	1	0.4	0.0	100±218
<i>Fusconaia flava</i>	32	13.6	0.0	3200±2177
<i>Lampsilis satura</i>	3	1.3	NV*	300±342
<i>Lampsilis teres</i>	1	0.4	0.0	100±218
<i>Leptodea fragilis</i>	11	4.7	NV	1100±883
<i>Megalonaia nervosa</i>	6	2.6	0.0	600±511
<i>Obliquaria reflexa</i>	1	0.4	0.0	100±218
<i>Plectomerus dombeyanus</i>	15	6.4	0.0	1500±977
<i>Pleurobema pyramidatum</i>	40	17.0	0.0	4000±2730
<i>Potamilus purpuratus</i>	6	2.6	0.0	600±511
<i>Quadrula metanevra</i>	1	0.4	0.0	100±218
<i>Quadrula pustulosa</i>	23	9.8	0.0	2300±1498
<i>Strophitus undulatus</i>	2	0.9	NV	200±295
<i>Tritogonia verrucosa</i>	10	4.2	0.0	1000±692
<i>Truncilla truncata</i>	7	3.0	NV	700±601
Totals	235	100.0	0.0	23400±9209

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 3M

Location: Mile 374.6

AR: Ouachita County

Stratum Size (n samples): 1.) 40m x 3m = 120m² (3)
 2.) 50m x 15m = 750m² (12)
 3.) 100m x 5m = 500m² (4)

Samples: 19

Substrate: Gravel

Minimum-Maximum density (#/m²): 1.) 4.0-8.0
 2.) 8.0-23.0
 3.) 14.0-50.0

Mean density #/m² (Standard deviation): 1.) 6.0 (2.3)
 2.) 13.7 (4.5)
 3.) 30.8 (17.9)
 Total: 16.0 (11.5)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	1	0.4	0.0	125±260
<i>Amblema plicata</i>	19	7.6	0.0	1307±620
<i>Ellipsaria lineolata</i>	15	6.0	0.0	1392±1003
<i>Elliptio dilatata</i>	14	5.6	0.0	1602±1352
<i>Fusconaia ebena</i>	7	2.8	0.0	813±1245
<i>Fusconaia flava</i>	35	14.1	0.0	6058±2085
<i>Lampsilis saturata</i>	1	0.4	NV*	63±409
<i>Ligumia recta</i>	1	0.4	0.0	125±260
<i>Megalonaia nervosa</i>	3	1.2	0.0	188±203
<i>Plectomerus dombeyanus</i>	8	3.2	0.0	500±644
<i>Pleurobema pyramidatum</i>	86	34.5	0.0	7142±2463
<i>Potamilus purpuratus</i>	4	1.6	0.0	313±319
<i>Quadrula pustulosa</i>	49	19.7	0.0	3808±2533
<i>Quadrula quadrula</i>	1	0.4	0.0	125±260
<i>Strophitus undulatus</i>	2	0.8	NV	200±295
<i>Tritogonia verrucosa</i>	1	0.4	0.0	125±260
<i>Truncilla donaciformis</i>	1	0.4	NV	125±260
<i>Truncilla truncata</i>	1	0.4	NV	125±260
Totals	249	99.9	0.0	26345±9514

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 5M				
Location: Mile 370.0				
AR: Ouachita County				
Stratum Size (n samples):	1.)	75m x 5m = 375m ²	(4)	
	2.)	75m x 10m = 750m ²	(7)	
Total Samples:	11			
Substrate:	Silty sand over gravel			
Minimum-Maximum density (#/m ²):	1.)	5.0-28.0		
	2.)	11.0-129.0		
Mean density #/m ² (Standard deviation):	1.)	13.5 (10.2)		
	2.)	50.1 (46.3)		
		Total: 36.8 (40.7)		
Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	14	3.5	0.0	1473±1867
<i>Amblema plicata</i>	11	2.7	0.0	1125±1000
<i>Cyprogenia aberti</i>	1	0.2	0.0	107±196
<i>Ellipsaria lineolata</i>	11	2.7	0.0	1179±702
<i>Elliptio dilatata</i>	12	3.0	0.0	1232±1004
<i>Fusconaia ebena</i>	5	1.2	0.0	536±469
<i>Fusconaia flava</i>	78	19.4	0.0	8009±5218
<i>Lampsilis abrupta</i>	1	0.2	ES*	94±205
<i>Lampsilis saturata</i>	1	0.2	NV**	107±232
<i>Leptodea fragilis</i>	2	0.5	NV	214±303
<i>Megalonaia nervosa</i>	1	0.2	0.0	107±232
<i>Plectomerus dombeyanus</i>	3	0.7	0.0	321±485
<i>Pleurobema coccineum</i>	20	5.0	0.0	2143±2691
<i>Pleurobema pyramidatum</i>	204	50.6	0.0	21629±17639
<i>Potamilus purpuratus</i>	2	0.5	0.0	214±278
<i>Quadrula metanevra</i>	3	0.7	0.0	321±485
<i>Quadrula pustulosa</i>	27	6.7	0.0	2893±1211
<i>Strophitus undulatus</i>	1	0.2	NV	107±232
<i>Truncilla truncata</i>	6	1.5	NV	643±908
Totals	403	99.7	0.0	42670±29039

** NV = No Commercial Value

* ES = Endangered Species

Appendix 1.3 cont'd

BED: 8M

Location: Mile 361.4

AR: Ouachita County

Stratum Size: 200m x 5m = 1000m²

Total Samples: 10

Substrate: Sand, gravel

Minimum-Maximum density (#/m²): 11.0-49.0Mean density #/m² (Standard deviation): 26.8 (12.5)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	3	1.1	0.0	300±343
<i>Amblema plicata</i>	32	12.0	0.0	3200±1703
<i>Ellipsaria lineolata</i>	21	7.8	0.0	2100±1030
<i>Elliptio dilatata</i>	5	1.9	0.0	500±918
<i>Fusconaia ebena</i>	1	0.4	0.0	100±125
<i>Fusconaia flava</i>	42	15.7	0.0	4200±2526
<i>Lampsilis saturata</i>	2	0.7	NV*	200±299
<i>Leptodea fragilis</i>	2	0.7	NV	200±299
<i>Megalonaia nervosa</i>	1	0.4	0.0	100±125
<i>Obliquaria reflexa</i>	5	1.9	0.0	500±503
<i>Plectomerus dombeyanus</i>	6	2.2	0.0	600±687
<i>Pleurobema pyramidatum</i>	52	19.4	0.0	5200±3669
<i>Potamilus purpuratus</i>	2	0.7	0.0	200±299
<i>Pyganodon grandis</i>	1	0.4	NV	100±125
<i>Quadrula metanevra</i>	1	0.4	0.0	100±125
<i>Quadrula nodulata</i>	5	1.9	0.0	500±604
<i>Quadrula pustulosa</i>	51	19.0	0.0	5100±2024
<i>Quadrula quadrula</i>	4	1.5	0.0	400±497
<i>Tritogonia verrucosa</i>	10	3.7	0.0	1000±1536
<i>Truncilla truncata</i>	22	8.2	NV	2200±1860
Totals	268	100.0	0.0	26800±8884

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 9M
 Location: Mile 361.1
 AR: Ouachita County
 Total Samples: 10
 Stratum Size: 100m x 6m = 600m²
 Substrate: Gravel, sand
 Minimum-Maximum density (#/m²): 9.0-57.0
 Mean density #/m² (Standard deviation): 27.1 (17.4)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	2	0.7	0.0	120±179
<i>Amblema plicata</i>	35	12.9	0.0	2100±1348
<i>Ellipsaria lineolata</i>	13	4.8	0.0	780±450
<i>Elliptio dilatata</i>	3	1.1	0.0	180±287
<i>Fusconaia ebena</i>	9	3.3	0.0	540±372
<i>Fusconaia flava</i>	62	22.9	0.0	3720±1676
<i>Lampsilis abrupta</i>	1	0.4	ES*	60±134
<i>Lampsilis hydiana</i>	1	0.4	NV**	60±134
<i>Lampsilis satula</i>	4	1.5	NV	240±297
<i>Leptodea fragilis</i>	1	0.4	NV	60±134
<i>Megalonaia nervosa</i>	2	0.7	0.0	420±450
<i>Obliquaria reflexa</i>	7	2.6	0.0	420±450
<i>Plectomerus dombeyanus</i>	10	3.7	0.0	600±448
<i>Pleurobema pyramidatum</i>	55	20.3	0.0	3300±3240
<i>Potamilus purpuratus</i>	4	1.5	50.0	240±297
<i>Ptychobranchus occidentalis</i>	4	1.5	NV	240±297
<i>Quadrula pustulosa</i>	36	13.3	0.0	2160±1362
<i>Quadrula quadrula</i>	2	0.7	0.0	120±179
<i>Tritogonia verrucosa</i>	14	5.2	0.0	840±944
<i>Truncilla truncata</i>	6	2.2	NV	360±411
Totals	271	100.1	0.7	16320±7397

* ES = Endangered Species

** NV = No Commercial Value

Appendix 1.3 cont'd

BED: 10M
 Location: Mile 360.3
 AR: Ouachita County
 Stratum Size: 150m x 18m = 2700m²
 Total Samples: 25
 Substrate: Gravel
 Minimum-Maximum density (#/m²): 7.0-48.0
 Mean density #/m² (Standard deviation): 22.4 (11.0)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	7	1.3	0.0	756±679
<i>Amblema plicata</i>	46	8.2	0.4	4968±1651
<i>Ellipsaria lineolata</i>	30	5.4	0.0	3240±1429
<i>Elliptio dilatata</i>	5	0.9	0.0	540±554
<i>Fusconaia ebena</i>	8	1.4	0.0	864±1046
<i>Fusconaia flava</i>	86	15.4	0.0	3288±4030
<i>Lampsilis satula</i>	6	1.0	NV*	648±579
<i>Lampsilis teres</i>	2	0.4	0.0	216±305
<i>Leptodea fragilis</i>	1	0.2	NV	108±221
<i>Megalonaia nervosa</i>	5	0.9	0.0	540±554
<i>Obliquaria reflexa</i>	3	0.5	0.0	324±486
<i>Plectomerus dombeyanus</i>	9	1.6	0.0	972±629
<i>Pleurobema pyramidatum</i>	184	32.9	0.0	19872±7763
<i>Potamilus purpuratus</i>	6	1.0	16.7	648±579
<i>Ptychobranchus occidentalis</i>	3	0.5	NV	324±486
<i>Pyganodon grandis</i>	1	0.2	NV	108±221
<i>Quadrula metanevra</i>	16	2.9	0.0	1738±1004
<i>Quadrula pustulosa</i>	84	15.0	0.0	9072±2494
<i>Quadrula quadrula</i>	2	0.4	0.0	108±221
<i>Tritogonia verrucosa</i>	15	2.7	0.0	1620±904
<i>Truncilla truncata</i>	41	7.3	NV	4428±1863
Totals	560	100.1	0.5	60480±12230

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 11M

Location: Mile 360.1

AR: Ouachita County

Stratum Size: 60m x 13m = 780m²

Total Samples: 10

Substrate: Sand, sand/gravel

Minimum-Maximum density (#/m²): 3.0-65.0Mean density #/m² (Standard deviation): 30.0 (21.1)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	6	2.0	0.0	468±467
<i>Amblema plicata</i>	22	7.3	4.5	1716±970
<i>Ellipsaria lineolata</i>	13	4.3	0.0	1014±741
<i>Elliptio dilatata</i>	8	2.7	0.0	624±817
<i>Fusconaia ebena</i>	31	10.3	0.0	2418±1779
<i>Fusconaia flava</i>	24	8.0	0.0	1872±2008
<i>Lampsilis abrupta</i>	1	0.3	ES*	78±175
<i>Megalonaia nervosa</i>	3	1.0	0.0	234±267
<i>Obliquaria reflexa</i>	4	1.3	0.0	312±387
<i>Plectomerus dombeyanus</i>	5	1.7	0.0	390±471
<i>Pleurobema pyramidatum</i>	118	39.3	0.0	9204±5806
<i>Potamilus purpuratus</i>	2	0.7	0.0	156±233
<i>Ptychobranchus occidentalis</i>	6	2.0	NV**	468±487
<i>Pyganodon grandis</i>	1	0.3	NV	78±175
<i>Quadrula metanevra</i>	8	2.7	0.0	624±681
<i>Quadrula pustulosa</i>	42	14.0	0.0	3267±2821
<i>Quadrula quadrula</i>	6	2.0	0.0	468±487
Totals	300	99.9	0.3	23712±910

* ES = Endangered Species

** NV = No Commercial Value

Appendix 1.3 cont'd

BED: 12M
 Location: Mile 359.7
 AR: Ouachita County
 Stratum Size: 100m x 11m = 1100m²
 Total Samples: 11
 Substrate: Sand, gravel, cobble
 Minimum-Maximum density (#/m²): 6.0-88.0
 Mean density #/m² (Standard deviation): 32.9 (22.6)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	5	1.1	0.0	500±603
<i>Amblema plicata</i>	19	5.2	0.0	1900±1278
<i>Ellipsaria lineolata</i>	8	2.2	0.0	800±579
<i>Elliptio dilatata</i>	4	1.1	0.0	400±371
<i>Fusconaia ebena</i>	15	4.1	0.0	1500±887
<i>Fusconaia flava</i>	39	10.8	0.0	3900±2426
<i>Lampsilis saturata</i>	1	0.3	NV*	100±221
<i>Megalonaia nervosa</i>	3	0.8	0.0	300±476
<i>Obliquaria reflexa</i>	3	0.8	0.0	300±344
<i>Plectomerus dombeyanus</i>	3	0.8	0.0	300±344
<i>Pleurobema pyramidatum</i>	193	53.3	0.0	19300±10454
<i>Potamilus purpuratus</i>	1	0.3	0.0	100±271
<i>Quadrula metanevra</i>	7	1.9	0.0	700±496
<i>Quadrula pustulosa</i>	33	9.1	0.0	3300±1316
<i>Quadrula quadrula</i>	4	1.1	0.0	100±271
<i>Tritogonia verrucosa</i>	5	1.4	0.0	500±496
<i>Truncilla truncata</i>	19	5.2	NV	1900±2160
Totals	362	99.5	0.0	36300±16619

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 13M

Location: Mile 359.3

AR: Ouachita County

Stratum Size (n samples): 1.) 80m x 7m = 560m² (5)
2.) 100m x 17m = 1700m² (17)

Total Samples: 22

Substrate: Sand, gravel

Minimum-Maximum density (#/m²): 1.) 8.0-16.0
2.) 6.0-70.0Mean density #/m² (Standard deviation): 1.) 12.0 (3.7)
2.) 27.1 (18.3)
Total: 23.3 (17.6)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	3	0.6	0.0	329±383
<i>Amblema plicata</i>	37	6.7	0.0	3677±1669
<i>Cyprogenia aberti</i>	1	0.2	0.0	94±193
<i>Ellipsaria lineolata</i>	27	5.1	0.0	2641±1145
<i>Elliptio dilatata</i>	11	2.1	0.0	1039±986
<i>Fusconaia ebena</i>	13	2.4	7.7	1319±1031
<i>Fusconaia flava</i>	39	7.3	0.0	3774±1734
<i>Lampsilis abrupta</i>	1	0.2	ES*	94±193
<i>Lasmigona costata</i>	1	0.2	NV**	94±193
<i>Megalonaia nervosa</i>	1	0.2	0.0	94±193
<i>Obliquaria reflexa</i>	9	1.7	0.0	941±649
<i>Plectomerus dombeyanus</i>	6	1.1	0.0	658±474
<i>Pleurobema pyramidatum</i>	198	37.1	0.0	19611±9383
<i>Potamilus purpuratus</i>	1	0.2	0.0	94±193
<i>Ptychobranchus occidentalis</i>	5	0.9	NV	472±207
<i>Quadrula metanevra</i>	14	2.6	0.0	1322±828
<i>Quadrula pustulosa</i>	101	18.9	0.0	9858±4165
<i>Quadrula quadrula</i>	5	0.9	0.0	472±207
<i>Tritogonia verrucosa</i>	13	2.4	0.0	1319±765
<i>Truncilla truncata</i>	49	9.2	NV	4764±1656
Totals	535	100.0	0.0	52714±15187

* ES = Endangered species

** NV = No Commercial Value

Appendix 1.3 cont'd

BED: 14M

Location: Mile 350.2

AR: Ouachita County

Stratum Size (n samples):
 1.) $80m \times 15m = 1200m^2$ (5)
 2.) $160m \times 15m = 2400m^2$ (6)
 3.) $185m \times 15m = 2775m^2$ (12)

Total Samples: 23

Substrate: Gravel, sand

Minimum-Maximum density (#/m²):
 1.) 18.0-57.0
 2.) 11.0-88.0
 3.) 6.0-114.0

Mean density #/m² (Standard deviation):
 1.) 33.8 (15.6)
 2.) 35.7 (27.3)
 3.) 34.2 (36.4)
 Total: 34.5 (29.6)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	25	3.1	8.0	5781±3957
<i>Amblema plicata</i>	26	3.3	0.0	7796±4797
<i>Cyprogenia aberti</i>	2	0.3	0.0	471±671
<i>Ellipsaria lineolata</i>	13	1.6	0.0	3548±2064
<i>Elliptio dilatata</i>	11	1.4	0.0	2553±2465
<i>Fusconaia ebena</i>	16	2.1	0.0	3700±2375
<i>Fusconaia flava</i>	92	11.6	0.0	25651±15953
<i>Lampsilis cardium</i>	6	0.8	NV*	1743±1524
<i>Leptodea fragilis</i>	1	0.1	NV	231±477
<i>Obliquaria reflexa</i>	31	3.9	0.0	8766±5457
<i>Plectomerus dombeyanus</i>	1	0.1	0.0	240±496
<i>Pleurobema pyramidatum</i>	164	20.7	0.0	44298±25172
<i>Ptychobranchus occidentalis</i>	11	1.4	NV	2925±1636
<i>Quadrula cylindrica</i>	2	0.3	0.0	800±1653
<i>Quadrula metanevra</i>	39	4.9	0.0	10741±4544
<i>Quadrula pustulosa</i>	157	19.8	0.0	46054±20236
<i>Quadrula quadrula</i>	16	2.0	0.0	4588±2302
<i>Strophitus undulatus</i>	2	0.3	NV	463±643
<i>Tritogonia verrucosa</i>	27	3.4	0.0	6749±4849
<i>Truncilla truncata</i>	151	19.0	NV	43831±13563
Totals	793	100.1	0.3	220973±81554

* NV = No Commercial Value

Appendix 1.3 cont'd

BED:	18M			
Location:	Mile 344.9			
AR:	Ouachita County			
Stratum Size (n samples):	1.) 100m x 8m = 800m ² (6)			
	2.) 170m x 13m = 2210m ² (19)			
Total Samples:	25			
Substrate:	Sand/gravel			
Minimum-Maximum density (#/m ²):	1.) 19.0-73.0			
	2.) 5.0-100.0			
Mean density #/m ² (Standard deviation):	1.) 48.5 (22.2)			
	2.) 46.0 (32.7)			
	Total: 46.6 (30.1)			
Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	2	0.2	0.0	233±1039
<i>Amblema plicata</i>	25	2.1	8.0	2993±1355
<i>Ellipsaria lineolata</i>	16	1.4	0.0	1929±1316
<i>Elliptio dilatata</i>	7	0.6	0.0	865±653
<i>Fusconaia ebena</i>	281	24.2	0.4	33859±11976
<i>Fusconaia flava</i>	179	15.4	0.0	21399±7186
<i>Lampsilis cardium</i>	2	0.2	NV*	250±355
<i>Leptodea fragilis</i>	1	0.1	NV	116±237
<i>Megalonaia nervosa</i>	30	2.6	0.0	3558±1732
<i>Obovaria reflexa</i>	3	0.3	0.0	349±389
<i>Plectomerus dombeyanus</i>	5	0.4	0.0	582±584
<i>Pleurobema pyramidatum</i>	460	39.6	0.0	55735±15817
<i>Ptychobranchus occidentalis</i>	6	0.5	NV	749±953
<i>Quadrula metanevra</i>	43	3.7	0.0	5206±2443
<i>Quadrula pustulosa</i>	79	6.8	0.0	9495±3483
<i>Quadrula quadrula</i>	6	0.5	0.0	732±734
<i>Tritogonia verrucosa</i>	3	0.3	0.0	349±389
<i>Truncilla donaciformis</i>	5	0.4	NV	582±470
<i>Truncilla truncata</i>	10	0.9	NV	1248±1149
Totals	1163	100.0	1.5	140060±36756

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 19M

Location: Mile 344.6

AR: Ouachita County

Stratum Size (n samples): 1.) 70m x 5m = 350m² (4)
2.) 120m x 12m = 1440m² (14)

Total Samples: 18

Substrate: Gravel, sand

Minimum-Maximum density (#/m²): 1.) 12.0-14.0
2.) 14.0-122.0Mean density #/m² (Standard deviation): 1.) 13.3 (1.0)
2.) 64.2 (30.8)
Total: 52.9 (34.7)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	2	0.2	0.0	306±291
<i>Amblema plicata</i>	27	2.8	0.0	2639±1062
<i>Ellipsaria lineolata</i>	8	0.8	0.0	792±758
<i>Elliptio dilatata</i>	8	0.8	0.0	823±1125
<i>Fusconaia ebena</i>	249	26.2	0.0	25350±7200
<i>Fusconaia flava</i>	142	14.9	0.0	14483±4553
<i>Lampsilis cardium</i>	3	0.3	NV*	293±458
<i>Megalonaia nervosa</i>	20	2.1	20.0	1904±1005
<i>Obliquaria reflexa</i>	2	0.2	0.0	206±291
<i>Plectomerus dombeyanus</i>	2	0.2	0.0	206±291
<i>Pleurobema coccineum</i>	2	0.2	0.0	191±274
<i>Pleurobema pyramidatum</i>	381	40.0	0.3	39127±13468
<i>Ptychobranchus occidentalis</i>	14	1.5	NV	1440±1577
<i>Quadrula cylindrica</i>	2	0.2	0.0	206±291
<i>Quadrula metanevra</i>	37	3.9	0.0	3806±2083
<i>Quadrula pustulosa</i>	40	4.2	0.0	4099±1906
<i>Quadrula quadrula</i>	4	0.4	0.0	411±491
<i>Strophitus undulatus</i>	1	0.1	NV	103±214
<i>Tritogonia verrucosa</i>	7	0.7	0.0	720±610
<i>Truncilla truncata</i>	1	0.1	NV	103±214
Totals	952	99.8	0.3	97106±24791

* NV = No Commercial Value

Appendix 1.3 cont'd.

BED: 20M

Location: Mile 344.5

AR: Ouachita County

Stratum Size (n samples): 1.) 300m x 20m = 6000m² (13)
2.) 250m x 20m = 5000m² (12)

Total Samples: 25

Substrate: Gravel, sand

Minimum-Maximum density (#/m²): 1.) 4.0-71.0
2.) 0.0-31.0Mean density #/m² (Standard deviation): 1.) 29.7 (25.8)
2.) 9.0 (11.2)
Total: 19.8 (22.4)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	3	0.6	0.0	1340±1514
<i>Amblema plicata</i>	16	3.2	0.0	7071±4685
<i>Ellipsaria lineolata</i>	6	1.2	0.0	2769±2657
<i>Elliptio dilatata</i>	1	0.2	0.0	462±944
<i>Fusconaia ebena</i>	100	20.2	0.0	45212±25952
<i>Fusconaia flava</i>	87	17.6	0.0	39571±20818
<i>Lampsilis cardium</i>	6	1.2	NV*	2724±3068
<i>Megalonaia nervosa</i>	12	2.4	0.0	5269±4460
<i>Obliquaria reflexa</i>	3	0.6	0.0	1340±1514
<i>Plectomerus dombeyanus</i>	3	0.6	0.0	1358±2049
<i>Pleurobema pyramidatum</i>	160	32.4	0.0	72500±34967
<i>Potamilus purpuratus</i>	3	0.6	0.0	1385±2049
<i>Ptychobranchus occidentalis</i>	7	1.4	NV	3186±3092
<i>Quadrula metanevra</i>	17	3.4	0.0	7667±2763
<i>Quadrula pustulosa</i>	42	8.5	0.0	18622±7728
<i>Quadrula quadrula</i>	8	1.6	0.0	3513±2488
<i>Strophitus undulatus</i>	2	0.4	NV	923±1286
<i>Tritogonia verrucosa</i>	4	0.8	0.0	3513±2488
<i>Truncilla donaciformis</i>	11	2.2	NV	5032±4514
<i>Truncilla truncata</i>	3	0.6	NV	1358±2049
Totals	494	99.7	0.0	223154±92718

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 21M

Location: Mile 340.9

AR: Ouachita County

Stratum Size (n samples): 1.) 70m x 20m = 1400m² (7)
 2.) 180m x 20m = 3600m² (19)

Total Samples: 26

Substrate: Gravel, sand

Minimum-Maximum density (#/m²): 1.) 8.0-28.0
 2.) 3.0-32.0

Mean density #/m² (Standard deviation): 1.) 21.0 (7.7)
 2.) 16.6 (8.7)
 Total: 17.8 (8.5)

Species	Number Collected of total	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	2	0.4	0.0	389±551
<i>Amblema plicata</i>	66	14.3	0.0	12663±4248
<i>Ellipsaria lineolata</i>	8	1.7	0.0	1547±1095
<i>Fusconaia ebena</i>	21	4.5	0.0	4021±1676
<i>Fusconaia flava</i>	55	11.9	0.0	10642±3192
<i>Lampsilis abrupta</i>	2	0.4	ES*	400±822
<i>Lampsilis hydiana</i>	1	0.2	NV**	189±391
<i>Lampsilis cardium</i>	10	2.2	NV	1926±1273
<i>Leptodea fragilis</i>	3	0.6	NV	568±850
<i>Megalonaia nervosa</i>	7	1.5	0.0	1347±1327
<i>Obliquaria reflexa</i>	16	3.5	0.0	3053±1327
<i>Obovaria</i> sp.	2	0.4	NV	389±551
<i>Plectomerus dombeyanus</i>	6	1.3	0.0	1147±1298
<i>Pleurobema pyramidatum</i>	84	18.1	0.0	16137±5221
<i>Potamilus purpuratus</i>	2	0.4	0.0	379±534
<i>Ptychobranchus occidentalis</i>	6	1.3	NV	1189±892
<i>Quadrula metanevra</i>	16	3.5	6.3	3074±1818
<i>Quadrula nodulata</i>	3	0.6	0.0	579±661
<i>Quadrula pustulosa</i>	75	16.2	0.0	14505±3707
<i>Quadrula quadrula</i>	58	12.5	0.0	11211±4646
<i>Strophitus undulatus</i>	4	0.9	NV	758±710
<i>Tritogonia verrucosa</i>	5	1.1	0.0	968±811
<i>Truncilla donaciformis</i>	1	0.2	NV	189±391
<i>Truncilla truncata</i>	10	2.2	NV	2000±855
Totals	463	99.9	0.2	105882±16608

* ES = Endangered Species

** NV = No Commercial Value

Appendix 1.3 cont'd

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	4	0.4	0.0	600±576
<i>Amblema plicata</i>	18	2.0	0.0	1800±1006
<i>Ellipsaria lineolata</i>	12	1.3	0.0	1800±1006
<i>Elliptio dilatata</i>	3	0.3	0.0	450±511
<i>Fusconaia ebena</i>	83	9.3	1.2	12450±3712
<i>Fusconaia flava</i>	157	17.5	0.6	23550±6373
<i>Lampsilis cardium</i>	16	1.8	NV*	1926±982
<i>Leptodea fragilis</i>	5	0.6	NV	750±629
<i>Megalonaia nervosa</i>	7	0.8	14.3	1050±1044
<i>Obovaria reflexa</i>	13	1.5	0.0	1950±902
<i>Obovaria</i> sp.	2	0.2	NV	300±427
<i>Plectomerus dombeyanus</i>	2	0.2	0.0	300±427
<i>Pleurobema pyramidatum</i>	321	35.8	0.0	48150±10958
<i>Potamilus purpuratus</i>	4	0.4	0.0	600±576
<i>Ptychobranchus occidentalis</i>	11	1.2	NV	1650±781
<i>Quadrula cylindrica</i>	3	0.3	0.0	450±676
<i>Quadrula metanevra</i>	46	5.1	0.0	6900±2541
<i>Quadrula nodulata</i>	1	0.1	0.0	150±308
<i>Quadrula pustulosa</i>	131	14.6	0.0	19650±4105
<i>Quadrula quadrula</i>	33	3.7	0.0	4950±2024
<i>Strophitus undulatus</i>	5	0.6	NV	750±770
<i>Tritogonia verrucosa</i>	1	0.1	0.0	150±304
<i>Truncilla truncata</i>	18	2.0	NV	2700±1509
Totals	896	99.8	0.3	134400±23383

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 23M

Location: Mile 339.5

AR: Ouachita/Calhoun County

Stratum Size (n samples): 1.) 150m x 25m = 3750m² (8)
2.) 175m x 25m = 4375m² (11)
3.) 75M x 25m = 1875m² (6)

Total Samples: 25

Substrate: Gravel/sand

Minimum-Maximum density (#/m²): 1.) 1.0-53.0
2.) 12.0-61.0
3.) 4.0-32.0Mean density #/m² (Standard deviation): 1.) 23.8 (22.2)
2.) 31.0 (15.4)
3.) 18.2 (11.5)

Total: 25.6 (17.3)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaia ligamentina</i>	11	1.7	9.1	4659±2029
<i>Amblema plicata</i>	44	6.9	0.0	17514±7194
<i>Ellipsaria lineolata</i>	20	3.1	0.0	7912±6651
<i>Elliptio dilatata</i>	16	2.5	0.0	6435±6505
<i>Fusconaia ebena</i>	51	8.0	0.0	21037±9721
<i>Fusconaia flava</i>	134	20.9	0.0	54290±29421
<i>Lampsilis cardium</i>	9	1.4	NV*	3551±6061
<i>Megalonaia nervosa</i>	18	2.8	16.7	71072±4049
<i>Obliquaria reflexa</i>	20	3.1	0.0	8026±2727
<i>Obovaria</i> sp.	1	0.2	0.0	398±819
<i>Plectomerus dombeyanus</i>	5	0.8	0.0	1818±1770
<i>Pleurobema pyramidatum</i>	145	22.7	0.0	59304±25043
<i>Potamilus purpuratus</i>	2	0.3	0.0	781±1137
<i>Ptychobranchus occidentalis</i>	2	0.3	NV	625±813
<i>Quadrula metanevra</i>	18	2.8	0.0	7074±8790
<i>Quadrula nodulata</i>	2	0.3	0.0	886±1241
<i>Quadrula pustulosa</i>	101	15.8	0.0	40525±13722
<i>Quadrula quadrula</i>	15	2.3	0.0	6435±4475
<i>Strophitus undulatus</i>	3	0.5	NV	1193±1755
<i>Tritogonia verrucosa</i>	11	1.7	0.0	4190±2657
<i>Truncilla truncata</i>	12	1.9	NV	5014±1694
Totals	640	100.0	0.6	258750±74347

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 24M
 Location: Mile 338.5
 AR: Ouachita/Calhoun County
 Stratum Size (n samples): 1.) 320m x 10m = 3200m² (17)
 2.) 140m x 10m = 1400m² (8)
 Total Samples: 25
 Substrate: Sandy gravel/cobble
 Minimum-Maximum density (#/m²): 1.) 4.0-59.0
 2.) 5.0-46.0
 Mean density #/m² (Standard deviation): 1.) 30.7 (16.1)
 2.) 16.8 (13.0)
 Total: 26.2 (16.4)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	6	0.9	16.7	1103±1047
<i>Amblema plicata</i>	65	10.0	0.0	12103±5227
<i>Ellipsaria lineolata</i>	11	1.7	0.0	2031±1104
<i>Elliptio dilatata</i>	5	0.8	0.0	941±750
<i>Fusconaia ebena</i>	27	4.1	0.0	5043±3691
<i>Fusconaia flava</i>	81	12.4	0.0	14982±6109
<i>Lampsilis cardium</i>	8	1.2	NV*	1435±1158
<i>Leptodea fragilis</i>	4	0.6	NV	753±697
<i>Megalonaia nervosa</i>	27	4.1	0.0	5056±3222
<i>Oboliquaria reflexa</i>	54	8.2	0.0	9993±3676
<i>Plectomerus dombeyanus</i>	16	2.4	12.5	2946±1713
<i>Pleurobema pyramidatum</i>	128	19.5	0.0	23737±9829
<i>Potamilus purpuratus</i>	11	1.7	0.0	2044±942
<i>Ptychobranchus occidentalis</i>	2	0.3	NV	363±518
<i>Quadrula metanevra</i>	7	1.1	0.0	1291±1029
<i>Quadrula nodulata</i>	11	1.7	0.0	2071±1865
<i>Quadrula pustulosa</i>	119	18.2	0.0	21976±5829
<i>Quadrula quadrula</i>	45	6.9	0.0	8418±2787
<i>Strophitus undulatus</i>	2	0.3	NV	376±773
<i>Tritogonia verrucosa</i>	20	3.0	5.0	3738±2754
<i>Truncilla truncata</i>	6	0.9	NV	1103±1127
Totals	655	100.0	0.8	121521±28364

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 25M
 Location: Mile 337.5
 AR: Ouachita County
 Stratum Size: 200m x 10m = 2000m²
 Total Samples: 20
 Substrate: Gravel, sand
 Minimum-Maximum density (#/m²): 10.0-37.0
 Mean density #/m² (Standard deviation): 19.1 (7.0)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	5	1.3	0.0	500±512
<i>Amblema plicata</i>	86	23.1	1.2	8600±2386
<i>Ellipsaria lineolata</i>	13	3.5	0.0	1300±954
<i>Elliptio dilatata</i>	1	0.3	0.0	100±208
<i>Fusconaia ebena</i>	19	5.1	0.0	1900±1553
<i>Fusconaia flava</i>	20	5.4	0.0	2000±1280
<i>Lampsilis cardium</i>	6	1.6	NV*	600±745
<i>Leptodea fragilis</i>	1	0.3	NV	100±208
<i>Megalonaia nervosa</i>	18	4.8	0.0	2000±1478
<i>Plectomerus dombeyanus</i>	6	1.6	16.7	600±745
<i>Pleurobema pyramidatum</i>	22	5.9	9.1	2200±949
<i>Potamilus purpuratus</i>	5	1.3	0.0	500±413
<i>Quadrula metanevra</i>	5	1.3	0.0	500±413
<i>Quadrula nodulata</i>	5	1.3	0.0	500±731
<i>Quadrula pustulosa</i>	77	20.7	0.0	7700±2223
<i>Quadrula quadrula</i>	53	14.2	0.0	5300±1576
<i>Tritogonia verrucosa</i>	6	1.6	0.0	600±437
<i>Truncilla truncata</i>	14	3.8	NV	1400±1005
Totals	372	97.1	1.0	38200±6499

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 26M
 Location: Mile 335.5
 AR: Calhoun County
 Stratum Size (n samples): 1.) 400m x 10m = 4000m² (17)
 2.) 200m x 10m = 2000m² (8)

Total Samples: 25

Substrate: Sandy gravel

Minimum-Maximum density (#/m²): 1.) 6.0-59.0
 2.) 2.0-78.0

Mean density #/m² (Standard deviation): 1.) 21.0 (12.7)
 2.) 45.9 (25.5)
 Total: 29.0 (20.9)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	9	1.2	11.1	2162±1558
<i>Amblema plicata</i>	73	10.1	2.7	17588±9401
<i>Ellipsaria lineolata</i>	22	3.0	0.0	5309±2619
<i>Elliptio dilatata</i>	3	0.4	0.0	706±1032
<i>Fusconaia ebena</i>	49	6.7	0.0	12015±5830
<i>Fusconaia flava</i>	137	18.9	0.0	33441±10710
<i>Lampsilis cardium</i>	7	1.0	NV*	1662±1302
<i>Leptodea fragilis</i>	2	0.3	NV	471±661
<i>Megalonaiaas nervosa</i>	31	4.3	0.0	7632±3756
<i>Obliquaria reflexa</i>	20	2.8	0.0	4765±2384
<i>Obovaria</i> sp.	1	0.1	NV	235±480
<i>Plectomerus dombeyanus</i>	6	0.8	0.0	1500±1289
<i>Pleurobema pyramidatum</i>	200	27.5	0.0	48809±17360
<i>Potamilus purpuratus</i>	2	0.3	50.0	471±661
<i>Quadrula metanevra</i>	19	2.6	0.0	4588±2647
<i>Quadrula nodulata</i>	1	0.1	0.0	235±1180
<i>Quadrula pustulosa</i>	95	13.1	0.0	22824±6628
<i>Quadrula quadrula</i>	35	4.8	0.0	8441±3423
<i>Tritogonia verrucosa</i>	4	0.1	0.0	956±938
<i>Truncilla donaciformis</i>	1	0.1	NV	235±480
<i>Truncilla truncata</i>	9	1.2	NV	2132±1124
Totals	726	99.4	0.6	176221±44009

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 27M
 Location: Mile 334.0
 AR: Ouachita County
 Stratum Size: 200m x 13m = 2600m²
 Total Samples: 25
 Substrate: Gravel, sand
 Minimum-Maximum density (#/m²): 4.0-65.0
 Mean density #/m² (Standard deviation): 23.0 (15.2)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	8	1.4	12.5	832±594
<i>Amblema plicata</i>	66	11.3	3.0	6864±2259
<i>Arkansas wheeleri</i>	1	0.2	ES*	104±213
<i>Ellipsaria lineolata</i>	10	1.7	0.0	1040±615
<i>Elliptio dilatata</i>	1	0.2	0.0	104±213
<i>Fusconaia ebena</i>	35	6.0	0.0	3640±2132
<i>Fusconaia flava</i>	84	14.4	0.0	8736±3587
<i>Lampsilis abrupta</i>	1	0.2	ES	104±213
<i>Lampsilis cardium</i>	6	1.0	NV**	624±465
<i>Leptodea fragilis</i>	2	0.3	NV	208±296
<i>Megalonaia nervosa</i>	31	5.3	0.0	3224±1279
<i>Obovaria reflexa</i>	15	2.6	0.0	1560±871
<i>Obovaria</i> sp.	2	0.3	NV	208±296
<i>Plectomerus dombeyanus</i>	13	2.2	0.0	1352±761
<i>Pleurobema pyramidatum</i>	123	21.1	0.0	12792±7044
<i>Potamilus purpuratus</i>	3	0.5	0.0	312±468
<i>Quadrula metanevra</i>	15	2.6	0.0	1560±1192
<i>Quadrula pustulosa</i>	117	20.0	0.0	12168±3209
<i>Quadrula quadrula</i>	32	5.5	0.0	3328±1427
<i>Strophitus undulatus</i>	3	0.5	NV	312±354
<i>Tritogonia verrucosa</i>	9	1.5	0.0	936±807
<i>Truncilla truncata</i>	7	1.2	NV	728±577
Totals	584	100.0	0.5	60736±16234

* ES = Endangered Species

** NV = No Commercial Value

Appendix 1.3 cont'd

BED: 28M

Location: Mile 331.8

AR: Calhoun County

Stratum Size (n samples): 1.) 75m x 6m = 450m² (5)
 2.) 70m x 15m = 1050m² (11)

Total Samples: 16

Substrate: 1.) Sandy gravel

2.) Cobble, gravel and sand

Minimum-Maximum density (#/m²): 1.) 9.0-18.0
 2.) 4.0-35.0

Mean density #/m² (Standard deviation): 1.) 12.4 (12.4)
 2.) 17.6 (8.6)
 Total: 16.0 (7.7)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	1	0.4	0.0	95±201
<i>Amblema plicata</i>	19	7.4	0.0	1770±1247
<i>Ellipsaria lineolata</i>	8	3.1	0.0	764±525
<i>Fusconaia ebena</i>	4	1.6	0.0	382±450
<i>Fusconaia flava</i>	17	6.6	0.0	1612±826
<i>Lampsilis cardium</i>	10	3.9	NV*	944±566
<i>Lampsilis teres</i>	1	0.4	0.0	90±190
<i>Megalonaia nervosa</i>	11	4.3	0.0	1045±1220
<i>Obovaria reflexa</i>	7	2.7	0.0	641±389
<i>Obovaria</i> sp.	1	0.4	NV	90±190
<i>Plectomerus dombeyanus</i>	9	3.5	0.0	815±592
<i>Pleurobema pyramidatum</i>	25	9.8	0.0	2375±1427
<i>Potamilus purpuratus</i>	2	0.8	0.0	191±270
<i>Ptychobranchus occidentalis</i>	1	0.4	NV	95±201
<i>Quadrula metanevra</i>	15	5.9	0.0	1432±1046
<i>Quadrula nodulata</i>	2	0.8	0.0	185±266
<i>Quadrula pustulosa</i>	89	34.8	0.0	8375±1980
<i>Quadrula quadrula</i>	18	7.0	0.0	1702±772
<i>Strophitus undulatus</i>	1	0.4	NV	90±190
<i>Tritogonia verrucosa</i>	7	2.7	0.0	652±488
<i>Truncilla truncata</i>	8	3.1	NV	753±500
Totals	256	100.0	0.0	24098±5993

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 29M

Location: 329.5

AR: Ouachita County

Stratum Size: 100m x 12m = 1200m²

Total Samples: 12

Substrate: Sand with pea gravel

Minimum-Maximum density (#/m²): 10.0-40.0Mean density #/m² (Standard deviation): 22.8 (10.8)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	1	0.4	0.0	100±218
<i>Amblema plicata</i>	27	9.9	0.0	2700±1943
<i>Ellipsaria lineolata</i>	3	1.1	0.0	300±471
<i>Fusconaia ebena</i>	6	2.2	0.0	600±605
<i>Fusconaia flava</i>	61	22.3	0.0	6100±3109
<i>Lampsilis cardium</i>	5	1.8	NV*	500±524
<i>Megalonaia nervosa</i>	9	3.3	0.0	900±571
<i>Obliquaria reflexa</i>	3	1.1	0.0	300±343
<i>Plectomerus dombeyanus</i>	11	4.0	0.0	100±755
<i>Pleurobema pyramidatum</i>	51	18.6	0.0	5100±3393
<i>Potamilus purpuratus</i>	4	1.4	25.0	400±373
<i>Quadrula metanevra</i>	4	1.4	0.0	400±494
<i>Quadrula pustulosa</i>	51	18.6	0.0	5100±2312
<i>Quadrula quadrula</i>	28	10.2	0.0	2800±1135
<i>Strophitus undulatus</i>	1	0.4	NV	100±218
<i>Tritogonia verrucosa</i>	9	3.3	0.0	900±471
Totals	274	100.0	0.4	27300±8160

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 30M
 Location: Mile 327.0
 AR: Calhoun County
 Stratum Size (n samples): 1.) 100m x 7m = 700m² (6)
 2.) 200m x 15m = 3000m² (19)
 Total Samples: 25
 Substrate: 1.) Sandy gravel
 2.) Pea gravel and sand
 Minimum-Maximum density (#/m²): 1.) 5.0-33.0
 2.) 0.0-41.0
 Mean density #/m² (Standard deviation): 1.) 20.2 (11.4)
 2.) 13.5 (9.6)
 Total: 15.1 (10.2)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	4	1.1	0.0	590±688
<i>Amblema plicata</i>	30	7.9	0.0	4325±1882
<i>Ellipsaria lineolata</i>	10	2.7	0.0	1496±900
<i>Elliptio dilatata</i>	2	0.5	0.0	275±900
<i>Fusconaia ebena</i>	6	1.6	0.0	906±900
<i>Fusconaia flava</i>	62	16.4	0.0	8470±4012
<i>Lampsilis cardium</i>	11	2.9	NV*	1496±900
<i>Lampsilis teres</i>	1	0.3	0.0	117±460
<i>Leptodea fragilis</i>	9	2.4	NV	1297±733
<i>Megalonaia nervosa</i>	31	8.2	0.0	4524±2609
<i>Obliquaria reflexa</i>	7	1.9	0.0	1064±925
<i>Plectomerus dombeyanus</i>	16	4.2	6.3	2320±1604
<i>Pleurobema pyramidatum</i>	20	5.3	0.0	2787±2011
<i>Potamilus purpuratus</i>	5	1.3	20.0	789±640
<i>Quadrula metanevra</i>	9	2.4	0.0	1297±1047
<i>Quadrula nodulata</i>	1	0.3	0.0	158±447
<i>Quadrula pustulosa</i>	88	23.3	0.0	12782±3147
<i>Quadrula quadrula</i>	35	9.3	0.0	5196±3187
<i>Strophitus undulatus</i>	3	0.8	NV	432±628
<i>Tritogonia verrucosa</i>	18	4.8	0.0	2718±1846
<i>Truncilla truncata</i>	9	2.4	NV	1380±2004
Totals	377	100.0	0.5	54538±15109

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 32M

Location: Mile 264.4

AR: Calhoun County

Stratum Size: 100m x 12m = 1200m²

Total Samples: 12

Substrate: Sandy clay with pea gravel

Minimum-Maximum density (#/m²): 1.0-29.0Mean density #/m² (Standard deviation): 11.3 (6.2)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	13	4.6	0.0	3120±1612
<i>Ellipsaria lineolata</i>	5	1.8	0.0	1200±1008
<i>Fusconaia ebena</i>	1	0.4	0.0	240±493
<i>Fusconaia flava</i>	7	2.5	0.0	1680±1335
<i>Lampsilis cardium</i>	10	3.6	NV*	2400±1884
<i>Leptodea fragilis</i>	7	2.5	NV	1680±1130
<i>Megalonaia nervosa</i>	28	10.0	0.0	6720±2599
<i>Obliquaria reflexa</i>	24	8.5	0.0	5760±2193
<i>Plectomerus dombeyanus</i>	25	8.9	0.0	6000±2251
<i>Pleurobema pyramidatum</i>	5	1.8	0.0	1200±1233
<i>Potamilus purpuratus</i>	2	0.7	0.0	480±685
<i>Quadrula metanevra</i>	4	1.4	0.0	960±923
<i>Quadrula nodulata</i>	2	0.7	0.0	480±685
<i>Quadrula pustulosa</i>	69	24.6	0.0	16560±5200
<i>Quadrula quadrula</i>	29	10.3	0.0	6960±2536
<i>Strophitus undulatus</i>	2	0.7	NV	480±685
<i>Tritogonia verrucosa</i>	3	1.1	0.0	720±818
<i>Truncilla truncata</i>	45	16.0	NV	10800±4669
Totals	281	100.1	0.4	67680±15144

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 36M

Location: Mile 271.8 Moro Bay State Park

AR: Calhoun County

Stratum Size (n samples): 1.) 200m x 15m = 3000m² (19)
2.) 50m x 15m = 750m² (6)

Total Samples: 25

Substrate: Sand with imbedded gravel

Minimum-Maximum density (#/m²): 1.) 14.0-58.0
2.) 20.0-93.0Mean density #/m² (Standard deviation): 1.) 29.0 (12.0)
2.) 67.6 (26.7)
Total: 38.2 (23.3)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	331	34.6	0.0	47592±8469
<i>Elliptio dilatata</i>	1	0.1	0.0	158±325
<i>Fusconaia ebena</i>	299	31.3	0.3	42803±14456
<i>Fusconaia flava</i>	38	4.0	0.0	5507±1671
<i>Lampsilis hydiana</i>	3	0.3	NV*	441±503
<i>Lampsilis teres</i>	2	0.2	0.0	316±445
<i>Leptodea fragilis</i>	3	0.3	NV	408±451
<i>Megalonaia nervosa</i>	14	1.5	0.0	1947±997
<i>Obliquaria reflexa</i>	12	1.3	0.0	1796±1133
<i>Plectomerus dombeyans</i>	125	13.1	2.4	17993±5828
<i>Pleurobema pyramidatum</i>	8	0.8	0.0	1000±1357
<i>Potamilus purpuratus</i>	9	0.9	22.2	1322±905
<i>Quadrula pustulosa</i>	71	7.4	0.0	10322±2174
<i>Quadrula quadrula</i>	20	2.1	0.0	3092±1479
<i>Tritogonia verrucosa</i>	5	0.5	0.0	789±764
<i>Truncilla truncata</i>	15	1.6	NV	2105±937
Totals	956	100.0	0.6	137592±23280

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 38M

Location: Mile 254.0 Storm Hole Bend

AR: Union County

Stratum Size (n samples): 1.) 225m x 18m = 4050m² (15)
 2.) 135m x 10m = 1350m² (5)
 3.) 150m x 10m = 1500m² (5)

Total Samples: 25

Substrate: 1.) Sandy gravel
 2.) Gravel, sand
 3.) Sandy gravel

Minimum-Maximum density (#/m²): 1.) 7.0-55.0
 2.) 6.0-59.0
 3.) 2.0-41.0

Mean density #/m² (Standard deviation): 1.) 25.7 (12.4)
 2.) 20.8 (22.1)
 3.) 15.0 (15.5)
 Total: 22.6 (15.2)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Actinonaias ligamentina</i>	1	0.2	0.0	270±557
<i>Amblema plicata</i>	183	32.4	0.0	50040±16507
<i>Fusconaia ebena</i>	139	24.6	0.0	38310±16117
<i>Fusconaia flava</i>	31	5.5	0.0	8520±4254
<i>Lampsilis cardium</i>	12	2.1	NV*	3360±1635
<i>Megalonaia nervosa</i>	14	2.5	0.0	3780±2738
<i>Obliquaria reflexa</i>	8	1.4	0.0	2190±1836
<i>Obovaria</i> sp.	7	1.2	0.0	1890±1968
<i>Plectomerus dombeyanus</i>	38	6.7	0.0	10380±4475
<i>Pleurobema pyramidatum</i>	4	0.7	0.0	1080±1717
<i>Potamilus purpuratus</i>	1	0.2	0.0	270±557
<i>Quadrula metanevra</i>	1	0.2	0.0	270±557
<i>Quadrula pustulosa</i>	85	15.1	0.0	23340±6569
<i>Quadrula quadrula</i>	16	2.8	0.0	4320±2603
<i>Tritogonia verrucosa</i>	3	0.5	0.0	810±889
<i>Truncilla truncata</i>	21	3.7	NV	5700±2976
Totals	564	99.8	0.0	197325±112275

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 39M

Location: Mile 252.4

AR: Union County

Stratum Size: 170m x 10m = 1700m²

Total Samples: 17

Substrate: Sandy gravel

Minimum-Maximum density (#/m²): 8.0-106.0Mean density #/m² (Standard deviation): 29.4 (28.8)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	149	29.8	0.0	19500±6968
<i>Fusconaia ebena</i>	163	32.6	0.0	16300±15303
<i>Fusconaia flava</i>	24	4.8	0.0	2400±2064
<i>Lampsilis hydiana</i>	2	0.4	NV*	200±288
<i>Lampsilis cardium</i>	16	3.2	NV	1600±719
<i>Leptodea fragilis</i>	11	2.2	NV	1100±700
<i>Megalonaia nervosa</i>	11	2.2	0.0	1100±527
<i>Obliquaria reflexa</i>	7	1.4	0.0	700±558
<i>Plectomerus dombeyanus</i>	27	5.4	0.0	2700±2018
<i>Pleurobema pyramidatum</i>	1	0.2	0.0	100±209
<i>Potamilus purpuratus</i>	4	0.8	25.0	400±422
<i>Quadrula pustulosa</i>	58	11.6	0.0	5800±1998
<i>Quadrula quadrula</i>	17	3.4	0.0	1700±870
<i>Tritogonia verrucosa</i>	3	0.6	0.0	300±341
<i>Truncilla truncata</i>	7	1.4	NV	700±558
Totals	500	100.0	0.2	50000±25044

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 40M

Location: Mile 252.0

AR: Bradley County

Stratum Size: 135m x 7m = 945m²

Total Samples: 10

Substrate: Sand and gravel

Minimum-Maximum density (#/m²): 11.0-37.0Mean density #/m² (Standard deviation): 21.8 (8.8)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	72	33.0	0.0	6804±2475
<i>Fusconaia ebena</i>	61	28.0	0.0	5765±3158
<i>Fusconaia flava</i>	6	2.8	0.0	567±566
<i>Lampsilis cardium</i>	4	1.8	NV*	378±474
<i>Leptodea fragilis</i>	3	1.4	NV	284±324
<i>Megalonaia nervosa</i>	9	4.1	0.0	851±864
<i>Obliquaria reflexa</i>	2	0.9	0.0	189±425
<i>Obovaria</i> sp.	2	0.9	0.0	189±425
<i>Potamilus purpuratus</i>	3	1.4	0.0	284±454
<i>Quadrula pustulosa</i>	37	17.0	0.0	3497±1144
<i>Quadrula quadrula</i>	15	6.9	0.0	1418±1014
<i>Tritogonia verrucosa</i>	1	0.5	0.0	95±212
<i>Truncilla donaciformis</i>	3	1.4	NV	284±324
Totals	218	100.1	0.0	21452±5891

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 41M				
Location: Mile 251.3				
AR: Union County				
Stratum Size: 1.) 75m x 12m = 900m ² (9)				
2.) 175m x 10m = 1750m ² (16)				
Total Samples: 25				
Substrate: Sandy gravel				
Minimum-Maximum density (#/m ²): 1.) 3.0-33.0				
2.) 7.0-28.0				
Mean density #/m ² (Standard deviation): 1.) 13.2 (9.2)				
2.) 16.3 (6.4)				
Total: 15.2 (7.5)				
Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	106	27.9	0.0	11191±4111
<i>Fusconaia ebena</i>	56	14.7	0.0	6050±2912
<i>Fusconaia flava</i>	10	2.6	0.0	1075±883
<i>Lampsilis hydiana</i>	3	0.8	NV*	319±362
<i>Lampsilis cardium</i>	14	3.7	NV	1513±739
<i>Leptodea fragilis</i>	2	0.5	NV	219±307
<i>Megalonaia nervosa</i>	21	5.5	0.0	2250±1095
<i>Obliquaria reflexa</i>	6	1.6	0.0	647±466
<i>Obovaria</i> sp.	1	0.3	0.0	109±225
<i>Plectomerus dombeyanus</i>	35	9.2	0.0	3631±2124
<i>Pleurobema pyramidatum</i>	2	0.5	0.0	209±299
<i>Potamilus purpuratus</i>	2	0.5	0.0	200±271
<i>Quadrula pustulosa</i>	83	21.8	0.0	8863±2346
<i>Quadrula quadrula</i>	14	3.7	0.0	1484±775
<i>Tritogonia verrucosa</i>	11	2.9	0.0	1166±882
<i>Truncilla donaciformis</i>	1	0.3	NV	109±225
<i>Truncilla truncata</i>	13	3.4	NV	1413±1025
Totals	380	99.9	0.0	40477±7891

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 42M
 Location: Mile 250.8
 AR: Bradley County
 Stratum Size (n samples): 1.) 50m x 10m = 500m² (5)
 2.) 150m x 13m = 1950m² (20)
 Total Samples: 25
 Substrate: 1.) Sand, clay, gravel
 2.) Silty sand, gravel
 Minimum-Maximum density (#/m²): 1.) 16.0-27.0
 2.) 8.0-54.0
 Mean density #/m² (Standard deviation): 1.) 22.6 (4.8)
 2.) 21.8 (12.5)
 Total: 21.9 (11.3)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	158	28.8	0.0	15470±3547
<i>Ellipsaria lineolata</i>	1	0.2	0.0	98±200
<i>Fusconaia ebena</i>	145	26.4	0.0	14250±6271
<i>Fusconaia flava</i>	34	6.2	0.0	3343±1443
<i>Lampsilis hydiana</i>	1	0.2	NV*	98±200
<i>Lampsilis cardium</i>	13	2.4	NV	1278±656
<i>Leptodea fragilis</i>	3	0.5	NV	295±337
<i>Megalonaia nervosa</i>	40	7.3	0.0	3928±1497
<i>Obliquaria reflexa</i>	1	0.2	0.0	98±200
<i>Plectomerus dombeyanus</i>	18	3.3	0.0	1768±842
<i>Pleurobema pyramidatum</i>	1	0.2	0.0	98±200
<i>Potamilus purpuratus</i>	1	0.2	0.0	100±205
<i>Quadrula metanevra</i>	1	0.2	0.0	98±200
<i>Quadrula pustulosa</i>	98	17.9	0.0	9570±2579
<i>Quadrula quadrula</i>	20	3.6	0.0	1955±1031
<i>Tritogonia verrucosa</i>	5	0.9	0.0	490±480
<i>Truncilla donaciformis</i>	1	0.2	NV	98±200
<i>Truncilla truncata</i>	7	1.3	NV	685±545
Totals	548	100.0	0.0	53318±11348

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 43M

Location: Mile 250.5

AR: Union County

Stratum Size: 100m x 12m = 1200m²

Total Samples: 12

Substrate: Sandy gravel

Minimum-Maximum density (#/m²): 8.0-39.0Mean density #/m² (Standard deviation): 26.7 (9.5)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	80	24.9	0.0	8000±3005
<i>Fusconaia ebena</i>	95	29.6	0.0	9500±3736
<i>Fusconaia flava</i>	9	2.8	0.0	900±657
<i>Lampsilis cardium</i>	11	3.4	NV*	1100±601
<i>Leptodea fragilis</i>	4	1.2	NV	400±373
<i>Megalonaia nervosa</i>	19	5.9	0.0	1900±1094
<i>Obliquaria reflexa</i>	5	1.6	0.0	500±390
<i>Obovaria</i> sp.	5	1.6	0.0	500±601
<i>Plectomerus dombeyanus</i>	10	3.1	0.0	1000±158
<i>Pleurobema pyramidatum</i>	1	0.3	0.0	100±218
<i>Potamilus purpuratus</i>	1	0.3	0.0	100±218
<i>Quadrula nodulata</i>	1	0.3	0.0	100±218
<i>Quadrula pustulosa</i>	62	19.3	0.0	6200±1795
<i>Quadrula quadrula</i>	12	3.7	0.0	1200±970
<i>Tritogonia verrucosa</i>	1	0.3	0.0	100±218
<i>Truncilla truncata</i>	5	1.6	NV	500±524
Totals	321	99.9	0.0	32100±7202

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 44M

Location: Mile 249.0 Caney Marais Bend

AR: Bradley County

Stratum Size (n samples): 1.) 130m x 15m = 1950m² (7)
2.) 200m x 25m = 5000m² (18)

Total Samples: 25

Substrate: 1.) Sandy gravel
2.) Pea gravel, sandMinimum-Maximum density (#/m²): 1.) 9.0- 19.0
2.) 11.0-142.0Mean density #/m² (Standard deviation): 1.) 14.4 (2.3)
2.) 34.7 (28.9)
Total: 29.0 (26.1)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	221	30.4	0.0	61416±33225
<i>Ellipsaria lineolata</i>	1	0.1	0.0	278±573
<i>Fusconaia ebena</i>	126	17.4	0.0	35005±18714
<i>Fusconaia flava</i>	52	7.2	0.0	14447±10672
<i>Lampsilis hydiana</i>	2	0.3	NV*	556±785
<i>Lampsilis cardium</i>	21	2.9	NV	5837±2914
<i>Lampsilis teres</i>	1	0.1	0.0	279±570
<i>Leptodea fragilis</i>	2	0.3	NV	556±785
<i>Megalonaia nervosa</i>	20	2.8	0.0	5560±2869
<i>Obovaria reflexa</i>	12	1.7	0.0	3335±2203
<i>Obovaria</i> sp.	5	0.7	0.0	1390±1738
<i>Plectomerus dombeyanus</i>	72	9.9	0.0	20009±5931
<i>Potamilus purpuratus</i>	6	0.8	0.0	1667±1230
<i>Quadrula metanevra</i>	1	0.1	0.0	278±573
<i>Quadrula pustulosa</i>	129	17.8	0.0	35848±8544
<i>Quadrula quadrula</i>	37	5.1	0.0	10285±6876
<i>Tritogonia verrucosa</i>	5	0.7	0.0	1390±1422
<i>Truncilla donaciformis</i>	5	0.7	NV	1389±1392
<i>Truncilla truncata</i>	8	1.1	NV	2225±1580
Totals	726	100.1	0.0	201747±70129

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 46M

Location: Mile 247.5

AR: Bradley County

Stratum Size: 165m x 5m = 825m²

Total Samples: 10

Substrate: sand, gravel, pea gravel, cobble

Minimum-Maximum density (#/m²): 8.0-32.0Mean density #/m² (Standard deviation): 19.2 (7.8)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	83	43.2	0.0	6848±2763
<i>Fusconaia ebena</i>	36	18.8	0.0	2970±3016
<i>Fusconaia flava</i>	5	2.6	0.0	413±414
<i>Lampsilis cardium</i>	14	7.3	NV*	1155±838
<i>Lampsilis teres</i>	1	0.5	0.0	83±185
<i>Megalonaia nervosa</i>	3	1.6	0.0	248±283
<i>Obliquaria reflexa</i>	1	0.5	0.0	83±185
<i>Plectomerus dombeyanus</i>	10	5.2	0.0	825±677
<i>Potamilus purpuratus</i>	2	1.0	0.0	165±371
<i>Quadrula pustulosa</i>	32	16.7	0.0	2640±820
<i>Quadrula quadrula</i>	3	1.6	0.0	248±283
<i>Truncilla donaciformis</i>	1	0.5	NV	83±185
<i>Truncilla truncata</i>	1	0.5	NV	83±185
Totals	192	100.0	0.0	15840±4580

* NV = No Commercial Value

Appendix 1.3 cont'd

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	107	35.2	0.0	10574±5348
<i>Fusconaia ebena</i>	23	7.6	0.0	2273±2784
<i>Fusconaia flava</i>	7	2.3	0.0	692±1255
<i>Lampsilis cardium</i>	19	6.3	NV*	1878±1090
<i>Lampsilis teres</i>	1	0.3	0.0	99±209
<i>Leptodea fragilis</i>	1	0.3	NV	99±209
<i>Megalonaia nervosa</i>	10	3.3	0.0	988±683
<i>Obliquaria reflexa</i>	15	4.9	0.0	1482±955
<i>Obovaria</i> sp.	1	0.3	0.0	99±209
<i>Plectomerus dombeyanus</i>	23	7.6	0.0	2273±1423
<i>Potamilus purpuratus</i>	1	0.3	0.0	99±209
<i>Quadrula nodulata</i>	2	0.7	0.0	198±285
<i>Quadrula pustulosa</i>	69	22.7	0.0	6819±3417
<i>Quadrula quadrula</i>	20	6.6	0.0	3377±576
<i>Tritogonia verrucosa</i>	5	1.6	0.0	494±847
Totals	304	100.0	0.0	30042±11486

* NV = No Commercial Value

Appendix 1.3 cont'd

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	153	33.0	0.0	14875±4331
<i>Arcidens confragosus</i>	1	0.2	NV*	97±203
<i>Fusconaia ebena</i>	63	13.6	0.0	6125±4593
<i>Fusconaia flava</i>	23	5.0	0.0	6125±4593
<i>Lampsilis cardium</i>	10	2.2	NV	972±798
<i>Lampsilis teres</i>	1	0.2	0.0	97±203
<i>Megalonaia nervosa</i>	32	6.9	0.0	3111±1313
<i>Obliquaria reflexa</i>	17	3.7	0.0	1653±1168
<i>Plectomerus dombeyanus</i>	50	10.8	0.0	4861±3233
<i>Potamilus purpuratus</i>	2	0.4	0.0	194±279
<i>Quadrula nodulata</i>	2	0.4	0.0	194±279
<i>Quadrula pustulosa</i>	66	14.2	0.0	6417±2202
<i>Quadrula quadrula</i>	31	6.7	0.0	3014±1573
<i>Tritogonia verrucosa</i>	4	0.9	0.0	389±474
<i>Truncilla donaciformis</i>	1	0.2	NV	97±203
<i>Truncilla truncata</i>	8	1.7	NV	778±678
Totals	464	100.1	0.0	45111±18021

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 49M

Location: Mile 245.5

AR: Union County

Stratum Size: 100m x 10m = 1000m²

Total Samples: 10

Substrate: Pea gravel over sand

Minimum-Maximum density (#/m²): 2.0-45.0Mean density #/m² (Standard deviation): 20.0 (13.2)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	42	21.0	0.0	4200±2918
<i>Fusconaia ebena</i>	40	20.0	0.0	4000±3073
<i>Fusconaia flava</i>	33	16.5	0.0	3300±2619
<i>Lampsilis hydiana</i>	1	0.5	NV*	100±225
<i>Lampsilis cardium</i>	5	2.5	NV	500±503
<i>Megalonaia nervosa</i>	7	3.5	0.0	700±824
<i>Obliquaria reflexa</i>	4	2.0	0.0	400±497
<i>Obovaria</i> sp.	1	0.5	0.0	100±225
<i>Plectomerus dombeyanus</i>	13	6.5	0.0	1300±1114
<i>Pleurobema pyramidatum</i>	1	0.5	0.0	100±225
<i>Quadrula pustulosa</i>	45	22.5	0.0	4500±1616
<i>Quadrula quadrula</i>	4	2.0	0.0	400±367
<i>Tritogonia verrucosa</i>	1	0.5	0.0	100±225
<i>Truncilla donaciformis</i>	1	0.5	NV	100±225
<i>Truncilla truncata</i>	2	1.0	NV	200±300
Totals	200	100.0	0.0	20000±9380

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 50M				
Location: Mile 245.4				
AR: Union County				
Stratum Size: 100m x 8m = 800m ²				
Total Samples: 10				
Substrate: Silty sand, gravel				
Minimum-Maximum density (#/m ²): 11.0-38.0				
Mean density #/m ² (Standard deviation): 21.5 (9.2)				
Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	78	36.3	0.0	6240±2755
<i>Fusconaia ebena</i>	15	7.0	0.0	1200±1144
<i>Fusconaia flava</i>	21	9.8	0.0	1680±1269
<i>Lampsilis cardium</i>	13	6.0	NV	1040±1969
<i>Megalonaia nervosa</i>	7	3.2	0.0	560±384
<i>Obliquaria reflexa</i>	3	1.4	0.0	240±274
<i>Plectomerus dombeyanus</i>	18	8.4	0.0	1440±958
<i>Pleurobema pyramidatum</i>	1	0.5	0.0	80±180
<i>Quadrula nodulata</i>	4	1.9	0.0	320±397
<i>Quadrula pustulosa</i>	35	16.3	0.0	2800±1424
<i>Quadrula quadrula</i>	16	7.4	0.0	1280±767
<i>Tritogonia verrucosa</i>	2	0.9	0.0	160±240
<i>Truncilla donaciformis</i>	1	0.5	NV	80±180
<i>Truncilla truncata</i>	1	0.5	NV	80±180
Totals	215	100.1	0.0	17280±5208

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 51M

Location: Mile 244.6

AR: Union County

Stratum Size (n samples): 1.) 70m x 7m = 490m² (6)
2.) 30m x 12m = 360m² (4)

Total Samples: 10

Substrate: 1.) Sand, gravel
2.) Pea gravel, sandMinimum-Maximum density (#/m²): 1.) 9.0-48.0
2.) 31.0-44.0Mean density #/m² (Standard deviation): 1.) 24.3 (15.7)
2.) 37.0 (7.0)
Total: 29.4 (14.0)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	94	32.0	0.0	8060±3870
<i>Fusconaia ebena</i>	39	13.3	0.0	3335±2850
<i>Fusconaia flava</i>	20	6.8	0.0	1700±1095
<i>Lampsilis cardium</i>	16	5.4	NV*	1365±876
<i>Leptodea fragilis</i>	1	0.3	NV	82±181
<i>Megalonaia nervosa</i>	13	4.4	0.0	1137±804
<i>Obovaria reflexa</i>	12	4.1	0.0	1022±460
<i>Obovaria</i> sp.	1	0.3	0.0	82±243
<i>Plectomerus dombeyanus</i>	21	7.1	0.0	1823±665
<i>Potamilus purpuratus</i>	1	0.3	0.0	82±181
<i>Quadrula pustulosa</i>	59	20.1	0.0	5102±1504
<i>Quadrula quadrula</i>	13	4.4	0.0	1103±393
<i>Tritogonia verrucosa</i>	3	1.0	0.0	262±421
<i>Truncilla truncata</i>	1	0.3	NV	90±200
Totals	294	99.8	0.0	25243±7259

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 52M
 Location: Mile 243.2
 AR: Bradley County
 Stratum Size (n samples): 1.) 180m x 4m = 720m² (7)
 2.) 100m x 15m = 1500m² (14)
 Total Samples: 21
 Substrate: 1.) Sand/gravel, pure sand
 2.) Sand/gravel, pure sand
 Minimum-Maximum density (#/m²): 1.) 6.0-50.0
 2.) 5.0-49.0
 Mean density #/m² (Standard deviation): 1.) 20.7 (17.2)
 2.) 19.2 (11.5)
 Total: 19.8 (13.3)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	130	31.2	0.0	13637±7256
<i>Arcidens confragosus</i>	1	0.2	NV*	103±213
<i>Fusconaia ebena</i>	41	10.0	0.0	4363±3152
<i>Fusconaia flava</i>	44	10.6	0.0	4659±4110
<i>Lampsilis cardium</i>	19	4.6	NV	2019±921
<i>Lampsilis teres</i>	1	0.2	0.0	107±220
<i>Leptodea fragilis</i>	1	0.2	NV	107±220
<i>Megalonaia nervosa</i>	18	4.3	0.0	1899±1231
<i>Obliquaria reflexa</i>	7	1.7	0.0	733±634
<i>Obovaria</i> sp.	3	0.7	0.0	317±518
<i>Plectomerus dombeyanus</i>	31	7.5	0.0	733±634
<i>Potamilus purpuratus</i>	1	0.2	0.0	107±220
<i>Quadrula nodulata</i>	7	1.7	0.0	746±568
<i>Quadrula pustulosa</i>	74	17.8	0.0	7873±2118
<i>Quadrula quadrula</i>	26	6.2	0.0	2721±1077
<i>Tritogonia verrucosa</i>	5	1.2	0.0	536±410
<i>Truncilla donaciformis</i>	2	0.5	NV	214±299
<i>Truncilla truncata</i>	5	1.2	NV	531±542
Totals	416	100.0	0.0	43736±13227

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 53M

Location: Mile 240.3

AR: Union County

Stratum Size (n samples): 1.) 150m x 15m = 2250m² (14)
 2.) 140m x 12m = 1680m² (11)

Total Samples: 25

Substrate: Silty sand with pea gravel

Minimum-Maximum density (#/m²): 1.) 10.0-36.0
 2.) 9.0-22.0

Mean density #/m² (Standard deviation): 1.) 20.9 (7.6)
 2.) 17.6 (3.9)
 Total: 19.4 (6.4)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	133	27.3	0.0	20912±4245
<i>Fusconaia ebena</i>	72	14.8	0.0	14410±5589
<i>Fusconaia flava</i>	47	9.7	0.0	7370±2239
<i>Lampsilis cardium</i>	25	5.1	NV*	3922±1397
<i>Leptodea fragilis</i>	2	0.4	NV	313±916
<i>Megalonaia nervosa</i>	19	3.9	0.0	2982±1252
<i>Obovaria reflexa</i>	15	3.1	0.0	2363±1326
<i>Obovaria</i> sp.	3	0.6	0.0	466±750
<i>Plectomerus dombeyanus</i>	29	6.0	0.0	4589±2028
<i>Pleurobema pyramidatum</i>	1	0.2	0.0	161±329
<i>Potamilus purpuratus</i>	1	0.2	100.0	153±314
<i>Quadrula nodulata</i>	6	1.2	0.0	964±856
<i>Quadrula pustulosa</i>	106	21.8	0.0	15929±4494
<i>Quadrula quadrula</i>	20	4.1	0.0	3182±2143
<i>Tritogonia verrucosa</i>	4	0.8	0.0	619±1563
<i>Truncilla donaciformis</i>	1	0.2	NV	161±329
<i>Truncilla truncata</i>	3	0.6	NV	474±537
Totals	487	100.0	0.2	76558±10188

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 54M

Location: Mile 239.1

AR: Union County

Stratum Size: 100m x 10m = 1000m²

Total Samples: 10

Substrate: Sand, clay, large gravel, cobble

Minimum-Maximum density (#/m²): 11.0-82.0Mean density #/m² (Standard deviation): 40.0 (23.9)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	100	25.0	0.0	10000±5629
<i>Fusconaia ebena</i>	135	33.7	0.0	13500±9211
<i>Fusconaia flava</i>	55	13.8	0.0	5500±3059
<i>Lampsilis cardium</i>	4	1.0	NV*	400±497
<i>Leptodea fragilis</i>	1	0.3	NV	100±125
<i>Megalonaia nervosa</i>	16	4.0	0.0	1600±1171
<i>Oliquaria reflexa</i>	2	0.5	0.0	200±300
<i>Plectomerus dombeyanus</i>	38	9.5	0.0	3800±1290
<i>Pleurobema pyramidatum</i>	1	0.2	0.0	100±125
<i>Potamilus purpuratus</i>	1	0.3	0.0	100±125
<i>Quadrula metanevra</i>	2	0.5	0.0	200±300
<i>Quadrula nodulata</i>	1	0.2	0.0	100±125
<i>Quadrula pustulosa</i>	34	8.5	0.0	3400±1811
<i>Quadrula quadrula</i>	9	2.2	0.0	900±623
<i>Tritogonia verrucosa</i>	1	0.3	0.0	100±125
 Totals	 400	 100.0	 0.0	 40000±16974

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 55M				
Location: Mile 236.0				
AR: Union County				
Stratum Size: 160m x 30m = 4800m ²				
Total Samples: 25				
Substrate: Sand, gravel, cobble				
Minimum-Maximum density (#/m ²): 6.0-46.0				
Mean density #/m ² (Standard deviation): 26.1 (11.0)				
Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	155	23.8	0.0	29760±6666
<i>Fusconaia ebena</i>	169	25.9	0.0	32448±10285
<i>Fusconaia flava</i>	21	3.2	0.0	4032±1909
<i>Lampsilis cardium</i>	6	0.9	NV*	1152±272
<i>Lampsilis teres</i>	1	0.2	0.0	192±394
<i>Leptodea fragilis</i>	8	1.2	NV	1536±940
<i>Megalonaia nervosa</i>	48	7.4	0.0	9216±3114
<i>Obliquaria reflexa</i>	8	1.2	0.0	1536±940
<i>Plectomerus dombeyanus</i>	88	13.5	0.0	16896±4968
<i>Potamilus purpuratus</i>	9	1.4	0.0	1728±1121
<i>Quadrula nodulata</i>	8	1.2	0.0	1536±1476
<i>Quadrula pustulosa</i>	84	12.9	0.0	16128±4551
<i>Quadrula quadrula</i>	34	5.2	0.0	6528±2602
<i>Tritogonia verrucosa</i>	5	0.8	0.0	960±986
<i>Truncilla donaciformis</i>	4	0.6	NV	768±738
<i>Truncilla truncata</i>	4	0.6	NV	768±738
Totals	652	100.0	0.0	125790±21652

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 56M

Location: Mile 233.5

AR: Ashley County

Stratum Size: 100m x 18m = 1800m²

Total Samples: 18

Substrate: Large gravel, sandy clay

Minimum-Maximum density (#/m²): 6.0-70.0Mean density #/m² (Standard deviation): 26.1 (14.5)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	142	30.3	0.0	14200±5954
<i>Fusconaia ebena</i>	131	27.9	0.0	13100±5101
<i>Fusconaia flava</i>	20	4.3	0.0	2000±1218
<i>Lampsilis cardium</i>	9	1.9	NV*	900±630
<i>Leptodea fragilis</i>	2	0.4	NV	200±289
<i>Megalonaia nervosa</i>	32	6.8	0.0	3200±1743
<i>Obovaria reflexa</i>	7	1.5	0.0	700±541
<i>Plectomerus dombeyanus</i>	41	8.7	0.0	4100±1776
<i>Potamilus purpuratus</i>	2	0.4	0.0	200±289
<i>Quadrula nodulata</i>	2	0.4	0.0	200±289
<i>Quadrula pustulosa</i>	62	13.2	0.0	6200±2569
<i>Quadrula quadrula</i>	16	3.4	0.0	1600±858
<i>Tritogonia verrucosa</i>	2	0.4	0.0	200±289
<i>Truncilla truncata</i>	1	0.2	NV	100±211
Totals	469	99.8	0.0	46900±12880

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 58M

Location: Mile 232.0

AR: Union County

Stratum Size (n samples): 1.) 100m x 8m = 800m² (8)
2.) 90m x 6m = 540m² (5)Substrate: 1.) Sand, gravel
2.) Gravel, sand

Total Samples: 13

Minimum-Maximum density (#/m²): 1.) 5.0-47.0
2.) 6.0-23.0Mean density #/m² (Standard deviation): 1.) 18.5 (14.6)
2.) 13.0 (6.4)
Total: 16.4 (12.1)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Ambloema plicata</i>	48	22.5	0.0	4680±2825
<i>Fusconaia ebena</i>	37	17.4	0.0	3570±2166
<i>Fusconaia flava</i>	4	1.9	0.0	400±563
<i>Lampsilis hydiana</i>	1	0.5	NV*	100±215
<i>Lampsilis cardium</i>	3	1.4	NV	300±452
<i>Leptodea fragilis</i>	2	0.9	NV	190±276
<i>Megalonaia nervosa</i>	9	4.2	0.0	870±633
<i>Obliquaria reflexa</i>	2	0.9	0.0	200±281
<i>Plectomerus dombeyanus</i>	66	31.0	0.0	6380±3706
<i>Pleurobema pyramidatum</i>	3	1.4	0.0	270±287
<i>Quadrula nodulata</i>	7	3.3	0.0	700±390
<i>Quadrula pustulosa</i>	10	4.7	0.0	960±622
<i>Quadrula quadrula</i>	18	8.5	0.0	1750±980
<i>Truncilla donaciformis</i>	2	0.9	NV	190±276
<i>Truncilla truncata</i>	1	0.5	NV	90±193
Totals	213	100.0	0.2	20650±9279

* NV = No Commercial Value

Appendix 1.3 cont'd

BED: 59M
 Location: Mile 230.9
 AR: Union County
 Stratum Size (n samples): 1.) 150m x 15m = 2250m² (11)
 2.) 150m x 20m = 3000m² (14)
 Substrate: 1.) Sand, embedded gravel
 2.) Sand, pea gravel
 Total Samples: 25
 Minimum-Maximum density (#/m²): 1.) 20.0-67.0
 2.) 6.0-65.0
 Mean density #/m² (Standard deviation): 1.) 37.4 (18.4)
 2.) 23.4 (14.9)
 Total: 29.7 (17.6)

Species	Number Collected	Percent of total	Percent legal	Population Estimate
<i>Amblema plicata</i>	233	31.4	0.0	48565±1886
<i>Fusconaia ebena</i>	195	26.3	0.5	42019±12715
<i>Fusconaia flava</i>	26	3.5	0.0	5464±3159
<i>Lampsilis cardium</i>	10	1.3	NV*	2133±1436
<i>Leptodea fragilis</i>	2	0.3	NV	419±596
<i>Megalonaia nervosa</i>	34	4.6	0.0	7032±2909
<i>Obliquaria reflexa</i>	30	4.0	0.0	6282±2223
<i>Plectomerus dombeyanus</i>	73	9.8	0.0	15282±5520
<i>Pleurobema pyramidatum</i>	2	0.3	0.0	419±596
<i>Potamilus purpuratus</i>	1	0.1	0.0	214±439
<i>Quadrula nodulata</i>	5	0.7	0.0	1062±863
<i>Quadrula pustulosa</i>	74	10.0	0.0	21886±4314
<i>Quadrula quadrula</i>	51	6.9	0.0	10865±3627
<i>Tritogonia verrucosa</i>	2	0.3	0.0	419±596
<i>Truncilla donaciformis</i>	1	0.1	NV	214±439
<i>Truncilla truncata</i>	3	0.4	NV	614±1261
Totals	742	100.0	0.1	154354±35536

* NV = No Commercial Value

APPENDIX 1.4 Physical parameters, species composition,
percent legal harvestable mussels and mean
densities for minor beds of the Ouachita
River, Arkansas, 1993-1995.

BED: 2m

Location: Mile 375.3

Stratum Size: Not available

Total Samples: 5

Substrate: Gravel

Minimum-Maximum density (#/m²): 8.0-15.0

Mean density #/m² (Standard deviation): 12.6 (2.9)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Actinonaias ligamentina</i>	3	4.8	0.0	0.6
<i>Amblema plicata</i>	2	3.2	0.0	0.4
<i>Cyprogenia aberti</i>	1	1.6	0.0	0.2
<i>Ellipsaria lineolata</i>	5	7.9	0.0	1.0
<i>Elliptio dilatata</i>	8	12.7	0.0	1.6
<i>Fusconaia flava</i>	8	12.7	0.0	1.6
<i>Leptodea fragilis</i>	2	3.2	NV*	0.4
<i>Plectomerus dombeyanus</i>	1	1.6	0.0	0.2
<i>Pleurobema pyramidatum</i>	6	9.5	0.0	1.2
<i>Potamilus purpuratus</i>	2	3.2	0.0	0.4
<i>Quadrula cylindrica</i>	1	1.6	0.0	0.2
<i>Quadrula metanevra</i>	1	1.6	0.0	0.2
<i>Quadrula pustulosa</i>	16	25.4	0.0	3.2
<i>Strophitus undulatus</i>	1	1.6	0.0	0.2
<i>Truncilla truncata</i>	6	9.5	NV	1.2
Totals	63	100.1	0.0	12.6

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 4m
 Location: Mile 370.5
 AR: Ouachita County
 Stratum Size: 70m x 3m = 210m²
 Total Samples: 5
 Substrate: Gravel
 Minimum-Maximum density (#/m²): 8.0-37.0
 Mean density #/m² (Standard deviation): 19.8 (12.9)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	15	15.2	0.0	3.0
<i>Ellipsaria lineolata</i>	1	1.0	0.0	0.2
<i>Elliptio dilatata</i>	5	5.0	0.0	1.0
<i>Fusconaia ebena</i>	1	1.0	0.0	0.2
<i>Fusconaia flava</i>	16	16.2	0.0	3.2
<i>Megalonaia nervosa</i>	2	2.0	0.0	0.4
<i>Plectomerus dombeyanus</i>	4	4.0	0.0	0.8
<i>Pleurobema pyramidatum</i>	50	50.5	0.0	10.0
<i>Quadrula pustulosa</i>	5	5.0	0.0	1.0
Totals	99	99.9	0.0	19.8

Appendix 1.4 cont'd

BED: 6m
 Location: Mile 367.6
 AR: Ouachita County
 Stratum Size: 85m x 3m = 255m²
 Total Samples: 5
 Substrate: Gravel
 Minimum-Maximum density (#/m²): 6.0-13.0
 Mean density #/m² (Standard deviation): 9.6 (2.8)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Actinonaia ligamentina</i>	2	4.2	0.0	0.4
<i>Amblema plicata</i>	10	20.8	0.0	2.0
<i>Elliptio dilatata</i>	1	2.1	0.0	0.2
<i>Fusconaia ebena</i>	1	2.1	0.0	0.2
<i>Fusconaia flava</i>	10	20.8	0.0	2.0
<i>Leptodea fragilis</i>	2	4.2	NV*	0.4
<i>Pleurobema pyramidatum</i>	14	29.2	0.0	2.8
<i>Quadrula nodulata</i>	1	2.1	0.0	0.2
<i>Quadrula pustulosa</i>	4	8.3	0.0	0.8
<i>Strophitus undulatus</i>	2	4.2	0.0	0.4
<i>Truncilla truncata</i>	1	2.1	NV	0.2
Totals	48	100.1	0.0	9.6

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 7m

Location: Mile 364.2

AR: Ouachita County

Stratum Size: 30m x 8m = 240m²

Total Samples: 5

Substrate: Gravel, fine sand

Minimum-Maximum density (#/m²): 15.0-49.0Mean density #/m² (Standard deviation): 23.0 (14.6)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Actinonaias ligamentina</i>	20	17.4	0.0	4.0
<i>Amblema plicata</i>	6	5.2	0.0	1.2
<i>Elliptio dilatata</i>	4	3.5	0.0	0.8
<i>Fusconaia ebena</i>	1	0.9	0.0	0.2
<i>Fusconaia flava</i>	30	26.1	0.0	6.0
<i>Pleurobema pyramidatum</i>	38	33.0	0.0	7.6
<i>Ptychobranchus occidentalis</i>	1	0.9	NV*	0.2
<i>Quadrula metanevra</i>	1	0.9	0.0	0.2
<i>Quadrula pustulosa</i>	14	12.2	0.0	2.8
Totals	115	100.1	0.0	23.0

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 15m

Location: Mile 351.7

AR: Calhoun County

Stratum Size: 40m x 10m = 400m²

Total Samples: 5

Substrate: Gravel

Minimum-Maximum density (#/m²): 10.0-37.0Mean density #/m² (Standard deviation): 22.2 (10.0)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Actinonaias ligamentina</i>	1	0.9	0.0	0.2
<i>Amblema plicata</i>	4	3.6	0.0	0.8
<i>Ellipsaria lineolata</i>	1	0.9	0.0	0.2
<i>Elliptio dilatata</i>	8	7.2	0.0	1.6
<i>Fusconaia ebena</i>	12	10.8	0.0	2.4
<i>Fusconaia flava</i>	25	22.5	0.0	5.0
<i>Megalonaia nervosa</i>	1	0.9	0.0	0.2
<i>Plectomerus dombeyanus</i>	3	2.7	0.0	0.6
<i>Pleurobema pyramidatum</i>	42	37.8	0.0	8.4
<i>Quadrula pustulosa</i>	10	9.0	0.0	2.0
<i>Strophitus undulatus</i>	1	0.9	NV*	0.2
<i>Tritogonia verrucosa</i>	1	0.9	0.0	0.2
<i>Truncilla truncata</i>	2	1.8	NV	0.4
Totals	111	99.9	0.0	22.2

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 16m

Location: Mile 351.5

AR: Calhoun County

Stratum Size: 40m x 12m = 480m²

Total Samples: 5

Substrate: Sand, pea gravel

Minimum-Maximum density (#/m²): 4.0-52.0Mean density #/m² (Standard deviation): 26.2 (21.2)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Actinonaias ligamentina</i>	1	0.8	0.0	0.2
<i>Amblema plicata</i>	5	3.8	0.0	1.0
<i>Ellipsaria lineolata</i>	10	7.6	0.0	2.0
<i>Elliptio dilatata</i>	4	3.1	0.0	0.8
<i>Fusconaia ebena</i>	20	15.3	0.0	4.0
<i>Fusconaia flava</i>	28	21.4	0.0	5.6
<i>Lampsilis abrupta</i>	1	0.8	ES*	0.2
<i>Megalonaia nervosa</i>	1	0.8	0.0	0.2
<i>Obovaria reflexa</i>	2	1.5	0.0	0.4
<i>Pleurobema pyramidatum</i>	28	21.4	0.0	5.6
<i>Quadrula metanevra</i>	6	4.6	0.0	1.2
<i>Quadrula pustulosa</i>	14	10.7	0.0	2.8
<i>Quadrula quadrula</i>	2	1.5	0.0	0.4
<i>Strophitus undulatus</i>	1	0.8	NV**	0.2
<i>Tritogonia verrucosa</i>	2	1.5	0.0	0.4
<i>Truncilla truncata</i>	6	4.6	NV	1.2
Totals	131	100.2	0.0	26.2

* ES = Endangered Species

** NV = No Commercial Value

Appendix 1.4 cont'd

BED: 17m

Location: Mile 350.1

AR: Ouachita County

Stratum Size: 50m x 8m = 400m²

Total Samples: 5

Substrate: Gravel

Minimum-Maximum density (#/m²): 25.0-122.0Mean density #/m² (Standard deviation): 62.2 (41.6)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Actinonaias ligamentina</i>	2	0.6	0.0	0.4
<i>Amblema plicata</i>	3	1.0	0.0	0.6
<i>Ellipsaria lineolata</i>	3	1.0	0.0	0.6
<i>Elliptio dilatata</i>	6	2.0	0.0	1.2
<i>Fusconaia ebena</i>	30	9.6	0.0	6.0
<i>Fusconaia flava</i>	63	20.3	0.0	12.6
<i>Lampsilis teres</i>	1	0.3	0.0	0.2
<i>Megalonaia nervosa</i>	2	0.6	0.0	0.4
<i>Obliquaria reflexa</i>	2	0.6	0.0	0.4
<i>Plectomerus dombeyanus</i>	3	1.0	0.0	0.6
<i>Pleurobema pyramidatum</i>	122	39.2	0.0	24.4
<i>Potamilus purpuratus</i>	1	0.3	0.0	0.2
<i>Ptychobranchus occidentalis</i>	5	1.6	NV*	1.0
<i>Quadrula metanevra</i>	11	3.5	0.0	2.2
<i>Quadrula pustulosa</i>	15	4.8	0.0	3.0
<i>Strophitus undulatus</i>	3	1.0	NV*	0.2
<i>Tritogonia verrucosa</i>	10	3.2	0.0	2.0
<i>Truncilla truncata</i>	29	9.3	NV	5.8
Totals	311	99.9	0.0	62.2

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 31m				
Location: Mile 321.0				
AR: Ouachita/Calhoun County				
Stratum Size: 50m x 2m = 100m ²				
Total Samples: 5				
Substrate: Clay, gravel, cobble				
Minimum-Maximum density (#/m ²): 7.0-22.0				
Mean density #/m ² (Standard deviation): 12.6 (6.3)				
Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Ambloema plicata</i>	9	14.3	0.0	1.8
<i>Fusconaia flava</i>	1	1.6	0.0	0.2
<i>Lampsilis cardium</i>	4	6.3	NV*	0.8
<i>Megalonaia nervosa</i>	10	15.9	0.0	2.0
<i>Obliquaria reflexa</i>	4	6.3	0.0	0.8
<i>Plectomerus dombeyanus</i>	2	3.2	0.0	0.4
<i>Potamilus purpuratus</i>	1	1.6	0.0	0.2
<i>Quadrula pustulosa</i>	10	15.9	0.0	2.0
<i>Quadrula quadrula</i>	21	33.3	0.0	4.2
<i>Truncilla truncata</i>	1	1.6	NV	0.2
Totals	63	100.0	0.0	12.6

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 33m
 Location: Mile 287.2
 AR: Union County
 Stratum Size: 75m x 5m = 375m²
 Total Samples: 5
 Substrate: Gravel, sand, clay
 Minimum-Maximum density (#/m²): 8.0-32.0
 Mean density #/m² (Standard deviation): 20.8 (8.9)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	33	31.7	0.0	6.6
<i>Ellipsaria lineolata</i>	1	1.0	0.0	0.2
<i>Fusconaia ebena</i>	1	1.0	0.0	0.2
<i>Lampsilis cardium</i>	3	2.9	NV*	0.2
<i>Megalonaia nervosa</i>	3	2.9	0.0	0.6
<i>Obovaria reflexa</i>	14	13.5	0.0	2.8
<i>Plectomerus dombeyanus</i>	1	1.0	0.0	0.2
<i>Quadrula nodulata</i>	3	2.9	0.0	0.8
<i>Quadrula pustulosa</i>	14	13.5	0.0	2.8
<i>Quadrula quadrula</i>	28	26.9	0.0	5.6
<i>Tritogonia verrucosa</i>	1	1.0	0.0	0.2
<i>Truncilla truncata</i>	2	1.9	NV	0.4
Totals	104	100.2	0.0	20.8

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 34m

Location: Mile 274.1

AR: Union County

Stratum Size: 50m x 3m = 150m²

Total Samples: 5

Substrate: Sand, gravel, cobble

Minimum-Maximum density (#/m²): 8.0-22.0

Mean density #/m² (Standard deviation): 17.6 (5.7)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	32	36.4	0.0	6.4
<i>Fusconaia flava</i>	1	1.1	0.0	0.2
<i>Megalonaia nervosa</i>	3	3.4	0.0	0.6
<i>Obliquaria reflexa</i>	4	4.5	0.0	0.8
<i>Plectomerus dombeyanus</i>	6	6.8	0.0	1.2
<i>Quadrula pustulosa</i>	20	22.7	0.0	4.0
<i>Quadrula quadrula</i>	19	21.6	0.0	3.8
<i>Tritogonia verrucosa</i>	1	1.1	0.0	0.2
<i>Truncilla truncata</i>	2	2.3	NV	0.4
Totals	88	99.9	0.0	17.6

Appendix 1.4 cont'd

BED: 35m

Location: Mile 272.4

AR: Union County

Stratum Size: 35m x 6m = 210m²

Total Samples: 5

Substrate: Coarse sand, pea gravel

Minimum-Maximum density (#/m²): 5.0-36.0Mean density #/m² (Standard deviation): 24.2 (12.7)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Actinonaias ligamentina</i>	1	0.8	0.0	0.2
<i>Amblema plicata</i>	84	69.4	0.0	16.8
<i>Arcidens confragosus</i>	1	0.8	NV*	0.2
<i>Fusconaia ebena</i>	3	2.5	0.0	0.6
<i>Fusconaia flava</i>	2	1.6	0.0	0.4
<i>Lampsilis cardium</i>	1	0.8	NV	0.2
<i>Lampsilis teres</i>	1	0.8	0.0	0.2
<i>Megalonaia nervosa</i>	6	5.0	0.0	1.2
<i>Obliquaria reflexa</i>	3	2.5	0.0	0.6
<i>Plectomerus dombeyanus</i>	2	1.6	0.0	0.4
<i>Potamilus purpuratus</i>	1	0.8	0.0	0.2
<i>Quadrula pustulosa</i>	7	5.8	0.0	1.4
<i>Quadrula quadrula</i>	6	5.0	0.0	1.2
<i>Truncilla truncata</i>	3	2.5	NV	0.6
Totals	121	99.9	0.0	24.2

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 37m

Location: Mile 270.4 Pigeon Hill Landing

AR: Union County

Stratum Size: 75m x 5m = 375m²

Total Samples: 5

Substrate: Sand, clay, gravel

Minimum-Maximum density (#/m²): 8.0-27.0Mean density #/m² (Standard deviation): 21.6 (7.8)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	55	50.9	0.0	11.0
<i>Fusconaia ebena</i>	4	3.7	0.0	0.8
<i>Lampsilis hydiana</i>	1	0.9	NV*	0.2
<i>Lampsilis cardium</i>	1	0.9	NV	0.2
<i>Megalonaia nervosa</i>	5	4.6	0.0	1.0
<i>Obliquaria reflexa</i>	3	2.8	0.0	0.6
<i>Plectomerus dombeyanus</i>	12	11.1	0.0	2.4
<i>Potamilus purpuratus</i>	1	0.9	0.0	0.2
<i>Quadrula pustulosa</i>	14	13.0	0.0	2.8
<i>Quadrula quadrula</i>	9	8.3	0.0	1.8
<i>Tritogonia verrucosa</i>	2	1.9	0.0	0.4
<i>Truncilla truncata</i>	1	0.9	NV	0.2
Totals	108	99.9	0.0	21.6

* NV = No Commercial Value

Appendix 1.4 cont'd

BED: 45m				
Location: Mile 248.0				
AR: Union County				
Stratum Size: 100m x 3m = 300m ²				
Total Samples: 5				
Substrate: Sandy gravel				
Minimum-Maximum density (#/m ²): 15.0-155.0				
Mean density #/m ² (Standard deviation): 75.6 (55.9)				
Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	167	44.2	0.0	33.4
<i>Ellipsaria lineolata</i>	1	0.3	0.0	0.2
<i>Fusconaia ebena</i>	103	27.2	0.0	20.6
<i>Fusconaia flava</i>	24	6.3	0.0	4.8
<i>Lampsilis cardium</i>	5	1.3	NV*	1.0
<i>Megalonaia nervosa</i>	14	3.7	0.0	2.8
<i>Obovaria reflexa</i>	4	1.0	0.0	0.8
<i>Obovaria</i> sp.	2	0.5	0.0	0.4
<i>Plectomerus dombeyanus</i>	14	3.7	0.0	2.8
<i>Pleurobema pyramidatum</i>	1	0.3	0.0	0.2
<i>Quadrula metanevra</i>	1	0.3	0.0	0.2
<i>Quadrula nodulata</i>	1	0.3	0.0	0.2
<i>Quadrula pustulosa</i>	23	6.1	0.0	4.6
<i>Quadrula quadrula</i>	12	3.2	0.0	2.4
<i>Tritogonia verrucosa</i>	1	0.3	0.0	0.2
<i>Truncilla donaciformis</i>	2	0.5	NV	0.4
<i>Truncilla truncata</i>	3	0.8	NV	0.6
Totals	378	100.0	0.0	75.6

* NV = No Commercial Value

Appendix 1.4 cont'd

BED:	57m			
Location:	Mile 233.0			
AR:	Ashley County			
Stratum Size:	80m x 4m = 320m ²			
Total Samples:	5			
Substrate:	Sand, pea gravel			
Minimum-Maximum density (#/m ²):	6.0-19.0			
Mean density #/m ² (Standard deviation):	12.0 (5.1)			
Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Aamblema plicata</i>	12	20.0	0.0	2.4
<i>Fusconaiia ebena</i>	10	16.7	0.0	2.0
<i>Megalonaiias nervosa</i>	2	3.3	0.0	0.4
<i>Obliquaria reflexa</i>	4	6.7	0.0	0.8
<i>Plectomerus dombeyanus</i>	6	10.0	0.0	1.2
<i>Potamilus purpuratus</i>	1	1.7	0.0	0.2
<i>Quadrula nodulata</i>	3	5.0	0.0	0.6
<i>Quadrula pustulosa</i>	18	30.0	0.0	3.6
<i>Quadrula quadrula</i>	4	6.7	0.0	0.8
Totals	60	100.1	0.0	26.5

Appendix 1.4 cont'd

BED: 60m

Location: Mile 223.0

AR: Ashley County

Stratum Size: 40m x 10m = 400m²

Total Samples: 5

Substrate: Sand, sand/gravel, sand/cobble

Minimum-Maximum density (#/m²): 7.0-29.0Mean density #/m² (Standard deviation): 19.8 (9.2)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	79	79.8	0.0	15.8
<i>Fusconaia ebena</i>	7	7.1	0.0	1.4
<i>Megalonaia nervosa</i>	2	2.0	0.0	0.4
<i>Obovaria</i> sp.	1	1.0	0.0	0.2
<i>Plectomerus dombeyanus</i>	3	3.0	0.0	0.6
<i>Quadrula pustulosa</i>	4	4.0	0.0	0.8
<i>Quadrula quadrula</i>	3	3.0	0.0	0.6
Totals	99	99.9	0.0	19.8

Appendix 1.4 cont'd

BED: 61m

Location: Mile 221.6

AR: Union County

Stratum Size: 40m x 10m = 400m²

Total Samples: 5

Substrate: Silt, sand

Minimum-Maximum density (#/m²): 10.0-22.0Mean density #/m² (Standard deviation): 16.4 (4.9)

Species	Number Collected	Percent of total	Percent legal	Mean Density
<i>Amblema plicata</i>	74	90.2	0.0	14.8
<i>Megalonaia nervosa</i>	1	1.2	0.0	0.2
<i>Plectomerus dombeyanus</i>	1	1.2	0.0	0.2
<i>Potamilus ohioensis</i>	1	1.2	0.0	0.2
<i>Quadrula pustulosa</i>	3	3.7	0.0	0.6
<i>Quadrula quadrula</i>	2	2.4	0.0	0.4
Totals	82	99.9	0.0	16.4